

From DEPARTMENT OF ONCOLOGY-PATHOLOGY  
Karolinska Institutet, Stockholm, Sweden

**Tumor-Associated Macrophages and  
Microglia:  
Double-edged sword in tumor evolvment and  
invasion**

Tatjana Wallmann



**Karolinska  
Institutet**

Stockholm 2018

All previously published papers were reproduced with permission from the publisher.

Published by Karolinska Institutet.

Printed by E-Print AB 2018

© Tatjana Wallmann, 2018

ISBN 978-91-7831-178-1

# **Tumor-Associated Macrophages and Microglia: Double-edged sword in tumor evolvement and invasion**

## **THESIS FOR DOCTORAL DEGREE (Ph.D.)**

Cancer Center Karolinska (CCK) Lecture Hall, R8:00, Karolinska University Hospital,  
Stockholm

**Friday, October 26<sup>th</sup>, 2018 at 12.30**

**Tatjana Wallmann**

*Principal Supervisor:*

Associate Professor Charlotte Rolny  
Karolinska Institutet  
Department of Oncology-Pathology

*Co-supervisor(s):*

Associate Professor Andreas Lundqvist  
Karolinska Institutet  
Department of Oncology-Pathology

Professor Arne Östman  
Karolinska Institutet  
Department of Oncology-Pathology

Associate Professor Ola Larsson  
Karolinska Institutet,  
Science for Life Laboratory  
Department of Oncology-Pathology

Associate Professor Tatiana Petrova  
University of Lausanne  
Department of Fundamental Oncology

*Opponent:*

Professor Claire Lewis  
The University of Sheffield  
Department of Oncology and Metabolism  
Division of Inflammation and Tumor Targeting

*Examination Board:*

Associate Professor Myriam Aouadi  
Karolinska Institutet  
Department of Medicine, Huddinge

Professor Lars-Gunnar Larsson  
Karolinska Institutet  
Department of Microbiology, Tumor and Cell  
Biology-Biomedicum

Associate Professor Lasse Jensen  
Linköping University  
Department of Medical and Health Sciences



“Run when you can,  
walk if you have to,  
crawl if you must;  
just never give up.”

(Dean Karnazes)

# **EIN KLEINER EINBLICK IN DIE KREBSENTSTEHUNG UND DIE ROLLE DER MAKROPHAGEN UND MIKROGLIA**

Krebs ist die zweithäufigste Todesursache weltweit und je älter wir werden, desto wahrscheinlicher ist es daran zu erkranken. Diese Krankheit macht vor Niemandem halt; ob Frau, ob Mann, ob alt, ob jung. Jeden kann es treffen und zu jedem Zeitpunkt im Leben. Am Anfang einer Behandlung stellt sich immer die Frage: „Kann ich den Kampf gegen den Krebs gewinnen?“ Und am Ende einer jeden Behandlung bleibt die Unsicherheit: „Habe ich den Kampf gegen den Krebs gewonnen?“ Zu Recht, denn es kann passieren, dass vereinzelte sehr resistente Krebszellen dem Tod entrinnen konnten und auf der Lauer liegen, damit sie später wieder zuschlagen können.....und diesmal stärker als zuvor. Ein Kampf, der zu häufig verloren wird.

Was aber ist Krebs? Einfach beschrieben sind es Zellen, die unkontrolliert wachsen. Was macht diese Krankheit dann so gefährlich?

Fangen wir von vorne an. Der menschliche Körper stellt ein System dar, in dem alle Prozesse nahezu perfekt aufeinander abgestimmt sind. In diesem System gibt es sehr viele verschiedene Kontrollinstanzen, die versichern, dass die Zellen funktionsfähig sind und ihre Aufgaben richtig erfüllen. Sollte das nicht mehr der Fall sein, kann unter anderem ein „Selbsttötungsprogramm“ eingeleitet werden. Dieser programmierte Zelltod ist in jeder Zelle gespeichert, so dass jede Zelle ihn durchlaufen kann, um sicher zu stellen, dass alle Abläufe im Körper weiterhin reibungslos verlaufen. Sei es die Eliminierung von überflüssigen Zellen während der embryonalen Entwicklung, ein Fehler in der DNA durch UV-Strahlung oder einfach nur die Erneuerung von Zellen; die Zelle weiß, wann sie zu gehen hat und Platz schaffen muss für eine neue gesunde Zelle. Die Entstehung einer neuen Zelle wird durch den Reparaturapparat streng kontrolliert, so dass sichergestellt wird, dass die DNA fehlerfrei ist und die Zelle ihre Erlaubnis bekommt zu leben.

Das Problem entsteht, wenn sich zu viele Fehler in der DNA summieren, die verschiedenen Kontrollinstanzen diese Fehler übersehen und sogar selber Defekte aufweisen. Das ist der Zeitpunkt, wo die Zellen dem Stoppsignal zum unkontrollierten Wachstum und sogar dem Selbsttötungsprogramm entgehen. Es bildet sich eine wuchernde und immer größer werdende Zellmasse, die sich unendlich oft teilen kann und unsterblich zu sein scheint. Diese Zellen schmeißen die ganzen kontrolliert ablaufenden Prozesse über einen Haufen....wachsen und

wachsen. Wenn die Zellmasse eine bestimmte Größe erreicht hat, braucht sie die Neubildung von Blutgefäßen um weiter wachsen zu können. Wie sonst soll sie Sauerstoff und Nährstoffe bekommen, die sie zum Überleben braucht? Wachstumsfaktoren werden im Übermaß ausgeschüttet, welche dazu führen, dass die neugebildeten Blutgefäße im Übermaß und unorganisiert wachsen. Diese Blutgefäße sind gekennzeichnet durch undichte Stellen und eine schlechte Durchblutung. Wenn nun diese Zellen in benachbartes Gewebe eindringen und durch die undichten Stellen in den Blutgefäßen zu entfernten Organen gelangen, werden sie zu bösartigen Krebszellen. Diese breiten sie sich unkontrolliert im Körper aus, sie streuen, und die Entstehung von neuen unkontrolliert wachsenden Zellmassen beginnt, den Metastasen. Krebs wird zu einer lebensbedrohlichen Krankheit.

Doch wo bleibt das Immunsystem? Das Immunsystem ist eins der elementarsten Funktionen im menschlichen Körper, das sehr komplex und auf einander abgestimmt ist. Es schützt uns vor gefährlichen Eindringlingen und stellt sicher, dass eigene entartete Zellen zerstört werden. Diese Aufgaben verlangen das Zusammenspiel von vielen verschiedenen Mitgliedern, wozu auch die Immunzellen zählen. Diese Immunzellen sind auch als weiße Blutkörperchen bekannt und unter ihnen befinden sich die Makrophagen. Makrophagen zirkulieren in unserem Körper und suchen die Umgebung nach Fremdkörpern ab. Treffen sie diese an, fressen sie die Fremdkörper auf, kommunizieren durch das Aussenden von vielen verschiedenen Stoffen mit ihrer Umgebung und suchen den Kontakt zu anderen Immunzellen auf. Diese und noch viele weitere initiierte Prozesse sorgen dafür, dass die Gefahr eliminiert wird. Darüber hinaus spielen sie auch eine wichtige Rolle in anderen Bereichen, die zum Erhalt des Gleichgewichts der Körperfunktionen beitragen. Diese sind unter anderem die Wundheilung und Reparatur von Gewebe, wo auch das Wachstum von neuen Blutgefäßen benötigt wird. Eine Fähigkeit, die Tumorzellen ausnutzen. Makrophagen befinden sich in allen Organen und selbst im Gehirn findet man sie. Dort sind sie als Mikroglia bekannt. Mikroglia erfüllen ganz spezielle und auf das Nervensystem abgestimmte Aufgaben.

Makrophagen und Mikroglia, die den Tumor infiltrieren, nennt man auch Tumor-assoziierte Makrophagen und Mikroglia (TAMMs). Je nachdem welche Signale sie aus ihrer Umgebung bekommen, können sie ein bestimmtes Erscheinungsbild annehmen. Durch ihre Plastizität können sie auch schnell zwischen unterschiedlichen Aktivierungszuständen wechseln und so viele verschiedene Aufgaben durchführen. Manche davon behindern das Krebswachstum und seine Streuung, sie sind anti-tumoral; und andere fördern diese Prozesse, sie sind pro-tumoral.

In den meisten Fällen eliminiert das Immunsystem (Makrophagen und Mikroglia sind hier mit eingeschlossen) entartete Krebszellen, die immer wieder versuchen das Immunsystem zu überlisten. Makrophagen und Mikroglia zeigen ein anti-tumorales Erscheinungsbild. Viele potenzielle Krebszellen fallen dem Immunsystem zum Opfer, jedoch nicht alle. Irgendwann schaffen es die Krebszellen das Immunsystem auszutricksen. Sie bilden Moleküle auf ihrer Oberfläche ab, die das Signal „Iss mich nicht“ vermitteln und senden Stoffe aus, die den Zerstörungsmodus der Immunzellen unterdrücken. Aus Feinden werden Verbündete. An Stelle den Krebs zu bekämpfen, helfen ihm nun Makrophagen und Mikroglia bei der Entstehung, beim Wachstum und bei der Streuung; sie sind pro-tumoral. Viele dieser Mechanismen sind schon erforscht, aber viele andere warten darauf ergründet zu werden. Das ist jedoch leichter gesagt als getan. Krebs ist eine sehr komplexe Krankheit, die in jedem Organ etwas anders aussieht und die wir bis zum heutigen Tag immer noch nicht komplett verstanden haben. Eine Krankheit, in der sich Makrophagen und Mikroglia als zweischneidiges Schwert erweisen können.

Heutzutage gibt es viele verschiedene Therapiemöglichkeiten und es werden immer mehr entwickelt. Und doch verlieren wir zu oft den Kampf gegen den Krebs. Es scheint zu viele Hürden zu geben, die es zu überwinden gibt. Man muss z.B. die Medikamente erfolgreich zu den Krebszellen befördern, obwohl die Blutgefäße um den Krebs herum durchlässig sind. Dann kommt hinzu, dass die Verabreichung eines Medikaments Krebszellen abtötet, die eine spezielle Mutation aufweisen. Jedoch bilden Krebszellen eine Resistenz gegenüber dem Medikament auf und es gibt genügend andere Krebszellen, die andere Mutationen besitzen. Und dann ist da noch das Immunsystem. Es gibt sehr viele verschiedene Ansätze das Immunsystem wieder „zurück zu programmieren“, so auch für die Makrophagen und Mikroglia. Diese Zellen so zu manipulieren, dass sie wieder anti-tumoral agieren. Eine sehr wichtige Erkenntnis bei alledem ist, dass eine Behandlungsmöglichkeit oft nicht ausreicht. Eine Kombination mit Einbezug des Immunsystems scheint vielversprechend zu sein. Jedoch ist der Weg zur Krebsbekämpfung noch lang. Um irgendwann den vollen Umfang an Behandlungsmöglichkeiten in der Krebstherapie nutzen zu können, brauchen wir das Wissen über jeden einzelnen Ablauf, der in Verbindung mit der Krebsentstehung, dem Krebswachstum und der Streuung steht. Dazu gehört auch die „Fehlfunktion“ des Immunsystems dem Krebs als Freund und nicht als Feind zu begegnen. Ich hoffe, dass ich mit meiner Doktorarbeit einen kleinen Teil dazu beigetragen habe, diese Krankheit besser zu verstehen, insbesondere welche Rolle Makrophagen und Mikroglia darin spielen können.







## ABSTRACT

Tumor-associated macrophages (TAMs) are among the most abundant cell types in the tumor microenvironment. TAM infiltration is usually linked to tumor progression, metastasis and poor clinical outcome in most human cancers. These cells sense a wide range of intra- and extracellular signals, such as chemokines and growth factors but also disturbances in pH or availability of oxygen. Due to their remarkable plasticity, TAMs can react immediately to these signals and acquire or switch to different phenotypes and activation states accordingly. For the past decade, TAMs have been recognized as a new therapeutic target. Even though diverse monotherapy strategies targeting TAMs have shown limited success, there are many different experimental studies that have shown promising results when approaching TAM subsets in combination with several other treatments, such as chemotherapies. However, our knowledge about the full scope of TAM heterogeneity and function in tumor evolution and invasion is still lacking.

In trying to improve the understanding about the versatile role of TAMs during tumor progression and invasion, several contributions have been made in this thesis. First, we show a novel mechanism whereby the TAM population can be skewed to contain mainly anti-tumoral M1-like Macrophages. Thus, we show that the overexpression of Semaphorin (SEMA)3A by tumor cells selectively induced the proliferation of M1-like macrophages and decreased the expansion of M2-like macrophages. This resulted in enhanced recruitment and activation of cytotoxic CD8<sup>+</sup> T lymphocytes and NK cells which in turn inhibited tumor growth. Second, we demonstrate that macrophage-derived Vascular Endothelial Growth Factor (VEGF)-C improved vessel functionality and thereby decreased pulmonary metastasis while tumor-derived VEGF-C increased vessel abnormalization and lung metastasis. Third, we elucidated that Cripto-1 vaccination decreased lung metastasis by inducing a humoral response leading to activated NK cell killing. Finally, we identified that microglia (resident macrophages of the brain) but not bone marrow-derived macrophages (BMDMs) induced Platelet-derived growth factor receptor (PDGFR)B expression on glioma cells and thereby enhanced their migratory capacity.



## LIST OF SCIENTIFIC PAPERS

- I. Wallerius M\*, **Wallmann T\***, Bartish M, Östling J, Mezheyeuski A, Tobin NP, Nygren E, Pangigadde P, Pellegrini P, Squadrito ML, Pontén F, Hartman J, Bergh J, De Milito A1, De Palma M, Östman A, Andersson J, Rolny C. **Guidance Molecule SEMA3A Restricts Tumor Growth by Differently Regulating the Proliferation of Tumor-Associated Macrophages**  
*Cancer Res. 2016 Jun 1;76(11):3166-78*
- II. **Wallmann T**, Landwehr LS, Squadrito ML, Wallerius M, Agardy D, Bartish M, Hartman J, Bergh J, De Palma M, Rolny C. **Macrophage-derived Vascular Endothelial Growth Factor C decreases hematogenous metastatic dissemination by normalizing the tumor vasculature**  
*Manuscript*
- III. Witt K\*, Ligtenberg MA\*, Conti L, Lanzardo S, Ruijter R, **Wallmann T**, Tufvesson-Stiller H, Rolny C, Lladser A, Lundqvist A, Cavallo F, Kiessling R. **Cripto-1 plasmid DNA vaccination targets metastasis and cancer stem cells in murine mammary carcinoma**  
*Cancer Immunol Res. 2018 Aug 24; in press*
- IV. **Wallmann T**, Zhang XM, Wallerius M, Bolin S, Joly AL, Sobocki C, Leiss L, Jiang Y, Bergh J, Holland EC, Enger P, Andersson J, Swartling F, Miletic H, Uhrbom L, Harris RA, Rolny C. **Microglia induce PDGFRB expression in glioma cells to enhance their migratory capacity**  
*Accepted, iScience*

\* *equal contribution*



# CONTENTS

1	Introduction.....	9
1.1	Breast Cancer.....	9
1.2	Glioma .....	10
1.3	The Immune System.....	11
1.4	Macrophages and Microglia.....	12
1.4.1	Origin and Maintenance.....	13
1.4.2	Classification and Activation .....	14
1.5	Tumor Immunology .....	16
1.5.1	Tumor Microenvironment .....	17
1.6	Tumor-Associated Macrophages and Microglia (TAMMs) .....	18
1.6.1	Recruitment.....	19
1.6.2	<i>In situ</i> Proliferation .....	20
1.6.3	Innate-adaptive crosstalk .....	22
1.6.4	Angiogenesis.....	23
1.6.5	Lymphangiogenesis .....	24
1.6.6	Invasion and Metastasis.....	25
1.7	Therapeutic Implications.....	26
2	Aims of the Thesis .....	28
3	Results and Discussion .....	29
3.1	Tumor mouse models .....	29
3.2	Study I .....	30
3.3	Study II.....	32
3.4	Study III.....	36
3.5	Study IV .....	38
4	Conclusions.....	41
5	Acknowledgements .....	43
6	References.....	48

## LIST OF ABBREVIATIONS

$\alpha$ -SMA	$\alpha$ -smooth muscle actin
bFGF	basic fibroblast growth factor
BMDM	Bone marrow-derived macrophage
CCL	Chemokine (C-C motif) ligand
CCR	Chemokine (C-C motif) receptor
CD	Cluster Differentiation
CNS	Central nervous system
COX2	Cytochrome c oxidase subunit 2
CSC	Cancer stem cell
CSF1	Colony stimulating factor 1
CSF1R	Colony stimulating factor 1 receptor
CX <sub>3</sub> CR1	CX <sub>3</sub> C chemokine receptor 1
CXCL	Chemokine (C-X-C motif) ligand
DC	Dendritic cell
ECM	Extracellular matrix
EGF	Epidermal growth factor
GAM	Glioma-associated microglia
GBM	Glioblastoma multiforme
GDNF	Glial-derived neurotrophic factor
HGF	Hepatocyte growth factor
HIF	Hypoxia-inducible factor
HSC	Hematopoietic stem cell
IBA1	Ionized calcium-binding adapter molecule 1
IDH	isocitrate dehydrogenase [NADP]
IFN	Interferon
IGF	Insulin-like growth factor
IL	Interleukin
IRF	Interferon regulatory factor
LEC	Lymphatic endothelial cell
MHC	Major histocompatibility complex



MMP	Matrix metalloproteinase
MMTV-PyMT	Mouse mammary tumor virus-polyoma middle T antigen
MRC	Mannose receptor C-type
NK cell	Natural killer cell
NO	Nitric oxide
NP	Neuropilin
PDAC	Pancreatic ductal adenocarcinoma
PDGF	Platelet-derived growth factor
PDGFR	Platelet-derived growth factor receptor
ROS	Reactive oxygen species
Sall	Sal-like
SEMA	Semaphorin
TAM	Tumor-associated macrophage
TAMM	Tumor-associated macrophage and microglia
TGF	Transforming growth factor
TNF	Tumor necrosis factor
uPA	Urokinase-type plasminogen activator
VEGF	Vascular endothelial growth factor
VEGFR	Vascular endothelial growth factor receptor




# 1 INTRODUCTION

Prior to 2011, there has been limited attention to the role of the tumor microenvironment in tumorigenesis. In 2000, Hanahan and Weinberg introduced six hallmarks of cancer that underlined the sequential progression of cells to cancer cells during multiple genetic alterations, which by then only involved the tumor cells (1). However, during the last decades it has become more and more evident that the tumor microenvironment plays an important role. Therefore, in 2011, Hanahan and Weinberg revised the hallmarks of cancer and added four additional hallmarks that include cancer-related inflammation and the active participation of tumor-associated stromal cells in cancer development (2).

## 1.1 BREAST CANCER

Breast cancer is the second most common type of cancer worldwide and the most common among women (3, 4). This disease is very heterogeneous and needs to be characterized precisely to choose the best treatment options (5). The most useful and important characteristics to predict prognosis and responsiveness to treatment have been proven to be the anatomical and histological determination (6). Anatomical staging is defined by the TNM categories, which are the tumor size (T), lymph node involvement (N) and the existence of metastases (M) (7). Histological features include the grading containing information about cell proliferation and the receptor status. Based on the receptor expression, breast cancer has been divided into four distinct molecular subtypes. These are luminal A, luminal B, HER2-enriched and basal-like breast cancers (**Figure 1**) (8).




**Figure 1. Molecular subtypes of breast cancer.**

Basal-like breast cancers overlap with triple-negative breast cancers (TNBC) but it is not the same disease. In fact, 20% of TNBC are not basal-like (9). TNBC resembles the most aggressive subtype of invasive ductal carcinoma, accounts for around 15-20% of breast cancer incidences and is associated with poor prognosis (10-12). TNBC shows higher metastatic potential compared to other subtypes and has the main metastatic sites in bone, lung, liver and brain (13, 14). Lymph node metastasis is observed less frequently but it is a crucial prognostic factor (15, 16). Breast cancer treatment is chosen based on the parameters described above and includes surgery, neoadjuvant and adjuvant therapy treatments (4).

## 1.2 GLIOMA

Gliomas are the most common primary tumors of the central nervous system (CNS) accounting for almost 80% of all malignant primary tumors (17). They include astrocytomas, oligodendrogliomas, ependymomas and mixed gliomas with astrocytoma being the most common type of gliomas (18). Gliomas can be classified into grades based on malignancy (I-IV), where grade IV is the most malignant, undifferentiated and invasive occurring type and denoted glioblastoma multiforme (GBM) (19). GBMs can arise from primary or secondary astrocytomas and together with diffuse (grade II) and anaplastic (grade III) astrocytomas most often occur in cerebral hemispheres (20). Primary GBMs arise *de novo*, while secondary GBMs progress slowly from low-grade astrocytomas (19). Grade II and III astrocytomas generally harbor a mutation in the cytoplasmic isocitrate dehydrogenase [NADP] (IDH)1, while GBMs are mostly IDH1 wildtype (20). Based on gene-expression analyses gliomas can be further subdivided into classical, mesenchymal, neural and proneural subtypes (**Figure 2**) (21). Low-grade gliomas are mostly of a neural and proneural subtype, while the classical and mesenchymal subtypes occur more frequently in high-grade gliomas (21, 22). The growth pattern of gliomas is characterized by tumor extensions and tumor microsatellites growing far away from the main tumor mass into the normal brain (23). Conventional therapy for GBMs such as surgery, radiation and chemotherapy prolongs survival but is not curative (24).



**Figure 2. Molecular subtypes of astrocytomas.** Adapted from (18).

### 1.3 THE IMMUNE SYSTEM

In a living organism, one of the immune system's primary functions is to resist pathogens. This host defense system is composed of the innate and adaptive immune systems. These subsystems complement each other and are highly balanced. The interplay between the innate and adaptive immune systems is essential to ensure long-lasting protection against infections and exhibits high complexity (25).

The innate immune system consists of the complement system, soluble factors such as cytokines, natural killer (NK) cells and phagocytic cells such as macrophages and dendritic cells (DCs) (26). Innate immunity enables the body to respond immediately against invading microorganisms and provides most of the every day protection. Since this protection is incomplete, the body requires adaptive immunity, which is a more sophisticated system. Adaptive immunity is tailor-made and antigen-specific against each intruder. Cells of the

adaptive immune system, B and T lymphocytes, possess unique antigen receptors, which enable them to efficiently eradicate pathogens. Although the adaptive immune response to primary infection by a pathogen is slower, the response is more powerful when the pathogen is encountered the second time due to long-lived memory B and T lymphocytes (27). This two-component immune system shows a complicated and fine-tuned network evolved to clear infections efficiently and to protect the host.


## **1.4 MACROPHAGES AND MICROGLIA**

Macrophages are leukocytes that are found in all tissues in differing abundances performing a variety of functions within and between distinct tissues. Although macrophages are mainly known for their phagocytic capacity, they are also involved in many other important processes that provide homeostasis, inflammation and immunity. In the different tissues, macrophages exert substantial functional specialization (28). Macrophages are highly plastic cells that adapt quickly to microenvironmental stimuli and have been shown to be able to respond effectively to environmental changes (29, 30).

In the brain, microglia are a unique population of long-lived tissue-resident cells that share numerous macrophage markers, such as colony stimulating factor 1 receptor (CSF1R), ionized calcium binding adaptor molecule (IBA)1 and F4/80 (31). As macrophages, microglia survey their surroundings and, if needed, phagocytose debris, apoptotic cells and pathogens. Microglia and macrophages regulate migration, differentiation and survival of different cell types. Depending on the activation signal they receive, macrophages and microglia produce a number of anti- and pro-inflammatory cytokines and growth factors (32). However, microglia differ in several aspects such as origin and gene transcription. Transcriptome analyses have revealed over 200 genes, such as Sal-like (Sall)1 and purigenic G-protein coupled P2y12 receptor, and several microRNAs that are uniquely expressed by microglia (31). This might not seem surprising since the brain has a unique microenvironment, where microglia exert very specific and distinct functions to maintain CNS integrity. Through these functions, microglia support processes such as neurogenesis, neuronal differentiation, migration and maintenance (33-35).

### 1.4.1 Origin and Maintenance

The origin of macrophages and microglia has been discussed for a long time. Lineage-tracing experiments in mice revealed that macrophages and microglia originate from diverse lineages. During embryonic development, macrophages derive first from the yolk sac followed by hematopoiesis of circulating monocytes in the fetal liver. After birth, hematopoiesis in the liver is replaced by bone marrow hematopoiesis, which is by then the major source of circulating monocytes (36). However, not all tissue-resident macrophages derive from circulating monocytes and persist in adulthood, as the skin, pancreas, liver, spleen and brain are populated by yolk sac progenitors, while in the kidneys and lung they have a chimaeric origin arising from yolk sac and bone marrow (Figure 3) (37, 38).



**Figure 3. Origin and ontogeny of tissue-resident macrophages and microglia.** Adapted from (38).

It is important to consider that both microglia and macrophages may reside in the CNS but develop independently from each other. In mice, at the eighth day after conception CD45<sup>-</sup> c-kit<sup>+</sup> erythromyeloid precursors are the earliest yolk sac progenitors that can give rise to microglia. Microgliogenesis is dependent on transcription factors Pu.1 and Interferon regulatory factor (Irf)8, and is modulated by matrix metalloproteinases (MMP)-8 and -9 (39). Meningeal macrophages derive from hematopoietic stem cells (HSCs) and turn over rapidly

(38). In comparison, microglia survive and renew themselves throughout lifetime without any influx of circulating monocytes under physiological conditions (40). Thus, depending on the tissue, the maintenance of tissue-specific macrophages and microglia ranges between being exclusively embryo-derived, like microglia in the brain, to being constantly replaced by bone marrow-derived macrophages (BMDMs), such as *lamina propria* macrophages, found in the intestine (**Figure 3**) (41, 42). Interestingly, monocytes that infiltrate tissues to replace embryonic macrophages by hematopoiesis, as observed in the heart, can adapt and differentiate into resident macrophages. This finding underlines that macrophages, independent of the origin, can exert similar functions when they receive tissue-specific signals (43).

#### 1.4.2 Classification and Activation


The classification of macrophages and microglia is challenging and many aspects have to be taken into account when describing distinct subtypes. It is important to consider the source of macrophages and microglia, the system used and the characterization of these cells using many different markers. Based on a framework including the phenotype, physiological activation and functional activity, macrophages have been classified into two subsets, the ‘classically activated’ M1 macrophages and the ‘alternatively activated’ M2 macrophages (44).

##### *Macrophages independent of origin*

In the presence of pathogens, necrosis or pro-inflammatory molecules such as interferon (IFN)- $\gamma$  or tumor necrosis factor (TNF)- $\alpha$ , macrophages adopt the M1-like phenotype. They release pro-inflammatory cytokines and chemokines (e.g. interleukin (IL)-1, IL-23, chemokine (C-C motif) ligand (CCL)8), are angiostatic, produce cytotoxic mediators, such as reactive oxygen species (ROS), and are involved in T helper (T<sub>H</sub>)1-mediated immune response (45-48). Their phagocytic capacity allows them to clear the body of debris, pathogens and cancer cells. Moreover, they express high levels of major histocompatibility complex (MHC) class II molecules that activate the adaptive immune system and recruit T lymphocytes by antigen presentation (49, 50). On the other hand, macrophages that respond to T<sub>H</sub>2 cytokines (IL-4, IL-13) acquire the M2-like phenotype. Thereby they down-regulate their pro-inflammatory activities and instead produce anti-inflammatory mediators such as IL-10 and transforming growth factor (TGF)- $\beta$  (51). M2-like macrophages harbor also high phagocytic capacity as they are involved in fibrosis, tissue repair and wound healing and



promote angiogenesis (52). Even if macrophages have acquired a specific phenotype, due to their remarkable plasticity, they react quickly to microenvironmental stimuli and switch between the different phenotypes to execute the tasks needed (**Figure 4**) (49).




**Figure 4. Summary of the main macrophage polarization states of activated macrophages.**  
Adapted and modified from (53).

### *Microglia*

Under homeostatic conditions microglia are known as ‘resting’ or ‘surveillant’ microglia that actively screen the CNS to maintain physiological functions (54). As soon as microglia encounter disturbances they become activated and adopt either a cytotoxic or neuroprotective profile depending on the signals they receive (55). Microglia have been divided into states of ‘classical activation’, ‘alternative activation’ and ‘acquired deactivation’ similar to that of extracranial macrophages (**Figure 5**) (56). In short, classical activated microglia respond to injury and infection producing pro-inflammatory cytokines such as IL-1 $\beta$ , TNF- $\alpha$ , nitric oxide (NO), ROS and proteases (57-59). Thereby they upregulate CD11b and IBA1 and start to express genes that are associated with antigen presentation such as MHC class II, cluster differentiation (CD)80 and CD86, resembling an M1-like phenotype of extracranial macrophages (60). Alternatively activated M2-like microglia respond to and release the cytokines IL-4 and IL-13 to promote anti-inflammation, tissue repair and the uptake of debris by upregulating scavenger receptors, such as Mannose receptor C-type (MRC)1, similarly to

extracranial macrophages (46, 61-63). Importantly, M2-microglia support neuron survival by releasing insulin-like growth factor (IGF)-1 (64).



**Figure 5. Summary of the main microglia polarization states.** Adapted and modified from (65).

As macrophages, microglia are very plastic cells that can switch between different phenotypes (66). Furthermore, the binary classification of macrophages and microglia into M1- and M2-like phenotypes is not as simplistic as described and includes many fine-tuned subtypes. For instance, macrophages and microglia can show expression of both M1- and M2-like markers that can be differently balanced depending on the tissue, signal and function (66, 67).


## 1.5 TUMOR IMMUNOLOGY

It is thought that the first immune response at the early stage of neoplastic progression resembles the response to acute tissue injury resulting in the infiltration of inflammatory cells (68). If the immune cells fail to eradicate the neoplasm, features of acute inflammation shift to features of chronic inflammation within the local tissue that fosters cancer development. During the persistent recruitment of immune cells, tumor cells secrete different soluble factors that alter the microenvironment and recruit additional myeloid cells and lymphocytes (69, 70). Once the immune cells arrive at the tumor site, they are exposed to many different

tumor secreted factors (e.g. CCL2, IL-4, IL-10, cytochrome c oxidase subunit (COX)2) that skew immune cells from an anti-tumoral phenotype to a pro-tumoral phenotype, so that tumor cells can evade immune destruction (71). As a consequence, these pro-tumoral immune cells release various molecules like growth factors, angiogenic factors, cytokines and proteases that promote tumor growth, invasion and metastatic dissemination (72-74).

### 1.5.1 Tumor microenvironment

The tumor resembles an autonomous organ that consists of many different components including tumor cells themselves, extracellular matrix (ECM), blood and lymphatic vessels and immune-inflammatory cells that represent the tumor microenvironment (TME) (75). The release of cytokines, growth factors and hormones by the different players adds another layer of complexity in a very dynamic network during tumor progression and metastasis (**Figure 6**). The number and heterogeneity of all factors varies in different tumor and cell types and therefore it is very important to consider these diversities in anti-cancer therapy (76, 77).



**Figure 6. Tumor microenvironment of solid tumors.** In the stroma different types of cells and molecules contribute to tumor progression and metastasis. Adapted and modified from (78).

## 1.6 TUMOR-ASSOCIATED MACROPHAGES AND MICROGLIA (TAMMS)

In the tumor microenvironment macrophages and microglia are known as tumor-associated macrophages and microglia (TAMMs) deriving from tissue-resident macrophages or microglia and from circulating monocytes of the peripheral blood (79). They infiltrate the tumor stroma and can represent the most abundant cell type with more than 50% of the tumor mass (80). In the majority of cancer types, except for prostate, gastric and colon cancer, high infiltration of TAMMs is associated with lower patient survival (81).

### *Macrophages*

Tumor-associated macrophages (TAMs) are major contributors to cancer-related inflammation and in a tumor TAMs consist of a heterogeneous population showing both M1- and M2-like phenotypes (as described above). During cancer initiation, accumulation of M1-like TAMs contributes to cancer elimination, known as immune surveillance. The accumulation of M1-like TAMs activates cytotoxic lymphocytes and thereby restricts tumor growth. However, tumor cells can escape the immune surveillance and secrete immunosuppressive factors skewing immune cells from an anti-tumoral to a pro-tumoral phenotype (82). Hence, the proportion and abundance of TAMs shifts toward an M2-like phenotype during tumor development (83, 84). This is important for tumor progression and metastasis as M2-like TAMs express high levels of angiogenic factors, anti-inflammatory cytokines and scavenging receptors (85). During the course of tumor growth, TAMs constantly undergo phenotypic changes in response to microenvironmental factors (86). In fact, depending on the niche, TAMs express different molecules and exert distinct functions. For instance, TAMs around blood vessels release pro-angiogenic factors like Vascular Endothelial Growth Factor (VEGF)-A to induce tumor angiogenesis, while TAMs at the invasive front produce MMP-2 and -9 to degrade ECM proteins and to facilitate tumor dissemination (87).

As mentioned earlier the M1/M2 polarization model is a very simplistic view on macrophage phenotypes. In 2014, transcriptome-based network analysis on human macrophages revealed a spectrum of different phenotypes upon stimulation of macrophages with diverse activation signals (88). Furthermore, the comparison of normal renal tissue with patient samples of clear cell renal cell carcinoma using single-cell analysis via mass cytometry identified 17 TAM phenotypes. In fact, our group performed single cell sequencing analyses of tumor-associated macrophages from a mouse mammary tumor virus-polyoma middle T antigen (MMTV-PyMT) tumor model and uncovered 9 different subgroups based on the transcriptome.

Intriguingly, these phenotypes change dynamically during tumor progression (unpublished data).

### *Microglia*

Glioma-associated microglia (GAMs) promote glioma growth, invasion, immunosuppression and angiogenesis by releasing cytokines, growth factors and enzymes similar to that of extracranial macrophages. Microglia within a brain tumor mass show also both M1- and M2-like phenotypes with the M2-like phenotype becoming the predominant one during tumor progression. In glioma patients, high infiltration of CD163<sup>+</sup> and CD204<sup>+</sup> (human M2-markers) microglia are associated to poor clinical prognosis (89, 90). The inhibition of CSF1R re-educates GAMs from an M2-like phenotype to an M1-like phenotype enhancing their phagocytic capacity and impairing tumor-promoting functions (91).

Even though TAMMs share many characteristics, it is also important to understand the differences, since the brain displays a unique microenvironment with special needs. Unless a neuropathological condition occurs, it is microglia that reside almost exclusively in the brain. The disruption of the blood brain barrier allows the infiltration of monocytes from the periphery (92). Comparing the expression profiles of TAMMs from glioma-bearing mice with naïve microglia, RNA microarray analysis has revealed a different expression profile for around 1,000 transcripts. These TAMMs showed only partial overlap in their expression pattern with known signatures from *in vitro* polarized M1- and M2-like macrophage subtypes (93).

#### **1.6.1 Recruitment**

As discussed before, tissue-resident macrophages can arise from distinct populations and monocytes can comprise one of this population. The CSF1/CSF1R axis plays an important role in monopoiesis and mice deficient in either one of the factors show decreased numbers of monocytes in the BM and circulation (94, 95). Immature monocytes are Ly6C<sup>hi</sup> and become upon maturation Ly6C<sup>lo</sup>, which can occur either in the BM or circulation (96, 97). Blood monocytes can be divided into two functional subsets, based on the expression of different markers, namely short-lived inflammatory CX<sub>3</sub>C chemokine receptor1<sup>lo</sup>Chemokine (C-C motif) receptor2<sup>+</sup> (CX<sub>3</sub>CR1<sup>lo</sup>CCR2<sup>+</sup>)Ly6C<sup>hi</sup> monocytes that circulate and are recruited to inflamed tissues and long-lived resident CX<sub>3</sub>CR1<sup>hi</sup>CCR2<sup>-</sup>Ly6C<sup>lo</sup> monocytes. These CX<sub>3</sub>CR1<sup>hi</sup>CCR2<sup>-</sup>Ly6C<sup>lo</sup> monocytes are the precursors for myeloid cells in the tissues under

homeostatic conditions and their development is dependent on the orphan nuclear hormone receptor transcription factor *Nr4a1* (98, 99). Expression of G-coupled receptor for sphingosine-1-phosphate (S1PR5) induces the release of patrolling Ly6C<sup>lo</sup> monocytes from the BM (100). These patrolling monocytes crawl along blood vessel walls scanning for damage and the presence of pathogens (99).

Under inflammatory conditions, such as during tumor growth, CCL2 is upregulated and recruits Ly6C<sup>hi</sup> monocytes. These inflammatory monocytes express MHC class II and are good antigen-presenting cells (101, 102). However, during tumor progression Ly6C<sup>hi</sup> monocytes can be the precursors for TAM subsets and differentiate into Ly6C<sup>int</sup> that give rise to MHC class II<sup>lo</sup> and MHC class II<sup>hi</sup> TAMs. Characterization of these two subsets revealed that MHC class II<sup>lo</sup> TAMs resemble the M2-like phenotype, are proangiogenic, enriched in hypoxia and efficiently suppress T cell proliferation. On the contrary, MHC class II<sup>hi</sup> TAMs express a repertoire of genes that resembles more an M1-like phenotype and are mainly found in normoxic regions (103). The infiltration, differentiation and proliferation of tumor-infiltrating Ly6C<sup>hi</sup> monocytes is also dependent on CSF1R signaling since blockage of this signaling pathway impairs these functions (104). Indeed, high levels of CSF1 are correlated with macrophage infiltration in human metastatic breast cancer (105). Homozygous null mutation of CSF1 in a mouse model of breast cancer reduced TAM infiltration and abolished tumor progression and dissemination. Overexpression, on the other hand, increased tumor growth and metastasis (106). Moreover, inhibition of the CSF1R did not only abrogate TAM infiltration but also increased the recruitment of CD8<sup>+</sup> T cells resulting in reduced mammary and cervical tumor growth (107).

### *Microglia*

Glioma cells secrete several cytokines and chemokines to recruit microglia into the tumor mass. For instance, CCL2 is also a major contributor to microglia infiltration as well as CX<sub>3</sub>CL1 that increases the ability for migration and adhesion (108, 109). Furthermore, growth factors like glial cell-derived neurotrophic factor (GDNF) and hepatocyte growth factor (HGF) have been identified as very potent chemoattractants (110, 111).

### **1.6.2 *In situ* Proliferation**

Although macrophages that are involved in inflammation and tumor progression are known to be recruited from BMDMs, recent evidence has implied that local proliferation of resident

macrophages and microglia also contributes to the expansion during pathological conditions (109, 112).

### *Macrophages*

Recent findings show that in an inflammatory context, such as atherosclerosis or during infection, local M2-like macrophage proliferation is a key event and leads to the expansion of this macrophage subtype (113). During nematode infection, controlled by CSF1, IL-4 drives local proliferation of macrophages indicating that proliferation is a feature of a T<sub>H</sub>2 inflammation-oriented response (114, 115). Intriguingly, proliferation and recruitment can be interlinked as observed in adipose tissue. Here, CCL2 induces macrophage proliferation but also monocyte entry into circulation and adipose tissue (116).

Macrophage proliferation has not only been observed under inflammatory conditions but can also be found in a tumor setting. For instance, in the spontaneous MMTVneu mouse model, that resembles HER2<sup>+</sup> human breast cancer, local proliferation of fully differentiated CD11b<sup>low</sup>F4/80<sup>high</sup> TAMs is the essential mechanism accounting for the accumulation of TAMs under CSF1R-dependent signaling. Monocyte depletion did not affect this TAM population (117). The presence of proliferating TAMs in human breast cancer material has been correlated to hormone receptor negative tumors, high grade and negative clinical outcome (118). Furthermore, we have shown that semaphorin (SEMA) 3A increases M1-like macrophage proliferation while it inhibits expansion of M2-like macrophages leading to a pro-inflammatory microenvironment that hampers tumor growth in mammary breast cancer (119). Interestingly, in a murine pancreatic ductal adenocarcinoma (PDAC) model both tissue-resident macrophages and inflammatory monocytes contribute to the local TAM population. These populations varied in the expression of cell surface markers and their function. While embryonically-derived MHC II<sup>low</sup> TAMs exhibited remodeling capacities and thereby fueled PDAC progression, monocyte-derived MHC II<sup>high</sup> TAMs played an important role in antigen presentation (120).

### *Microglia*


In different brain malignancies, depending on the pathological condition, various observations have been made regarding microglia proliferation and recruitment of BMDMs. For example, acute motor nerve damage triggers microglia expansion but not an influx of monocytes (121). On the contrary, in brain tumors, both tissue-resident microglia and recruited BMDMs are present. These recruited BMDMs, upon entry, can upregulate 'microglia-specific' genes, such as *Cx3cr1*. Therefore, both populations contribute to the pool of glioma-residing myeloid cells (92, 122). However, glioma cells secrete different factors

that were shown to trigger local microglia proliferation, such as CCL2, granulocyte-macrophage (GM)-CSF, M-CSF and VEGF-A (108, 123, 124). Accordingly, M-CSF/M-CSF1R paracrine communication between microglia and GBM augmented infiltration of microglia and GBM invasion *in vivo* (125).

### 1.6.3. Innate-adaptive crosstalk

#### *Macrophages and Microglia*

As mentioned earlier, at the early stage of tumor initiation, M1-like TAMMs promote anti-tumor responses by secreting pro-inflammatory factors such as TNF- $\alpha$  and chemokines like (C-X-C motif) ligand (CXCL)9 and CXCL10. The release of these chemokines mediates activation and recruitment of cytotoxic T lymphocytes and NK cells that show anti-tumoral properties (126). However, M2-like TAMMs lose these anti-tumoral activities (52, 127, 128). They secrete immune-suppressive molecules such as IL-10 and TGF- $\beta$  that down-regulate anti-tumoral cytotoxic T lymphocyte and NK cell activity (129, 130). Instead, these TAMMs stimulate the differentiation of naïve CD4<sup>+</sup> T lymphocytes into T<sub>H</sub>2 cells and regulatory T lymphocytes that augment anti-tumoral activity (**Figure 7**) (131, 132).




**Figure 7. Cross-talk between TAMMs and T cells.** M1-TAMMs activate CD8<sup>+</sup> T lymphocytes, while M2-TAMMs inhibit CD8<sup>+</sup> T lymphocytes and activate regulatory CD4<sup>+</sup> T lymphocytes.



### 1.6.4 Angiogenesis

Tumor angiogenesis, triggered by the angiogenic switch, induces the growth of new blood vessels from pre-existing vasculature to supply the tumor with oxygen, nutrients and is required for the tumor to grow beyond a size of 1-2 mm<sup>3</sup> (133-136). Hence, the balance between anti- and pro-angiogenic factors shifts toward a pro-angiogenic microenvironment (137). Many different factors regulating the angiogenic switch have been reported including VEGF-A, TGF- $\beta$  and ROS (138-140). In fact, hypoxia is a key driver of angiogenesis and the secretion of VEGF-A by cancer cells and recruited leukocytes triggers the development of a vascular network (135). These tumor blood vessels are characterized by a disorganized establishment, excessive vessel branching and enlarged vessels (141, 142). The discontinuous lining of endothelial cells and pericytes but also defective basement membranes result in leaky and poorly perfused vessels that increase tumor intravasation and hamper drug delivery (143, 144).

TAMMs are key effectors in angiogenesis and are known to accumulate in poorly vascularized necrotic areas and to surround blood vessels (145). They switch into the pro-angiogenic phenotype in the presence of stress factors such as low oxygen, low pH and high lactate concentration (146). Hypoxia appears to be one of the strongest factors promoting TAMM recruitment (147, 148). Once they have reached the hypoxic sites, hypoxia changes gene expression toward pro-angiogenic genes through the modulation of hypoxia-inducible factor (HIF)-1 and HIF-2 and other transcription factors (149, 150). In response, TAMMs secrete a number of pro-angiogenic growth factors including VEGF-A, CXCL2, TNF- $\alpha$  and basic fibroblast growth factor (bFGF) but also matrix modulating mediators such as MMPs, urokinase-type plasminogen activator (uPA) and its receptor uPAR (87, 148, 151, 152). For instance, MMP-9 expression by microglia increases glioma growth and invasion (153). Moreover, inhibition of MMP-9 activity and expression by infiltrating macrophages reduces the release of VEGF-A from the ECM and thereby inhibits angiogenesis and tumor growth in cervical cancer (154). Interestingly, hypoxia does not seem to drive the M2-phenotype but rather fine tunes M2-like TAMMs (155). TAMs that surround blood vessels and promote angiogenesis were identified to express the Tie2 receptor (156). Accumulation of Tie2<sup>+</sup> macrophages is correlated with microvascular density and distant metastasis (**Figure 8**) (157, 158).



**Figure 8. Modulatory functions of macrophages in the tumor microenvironment.** Adapted from Macrophages: Biology and Role in the Pathology of Diseases, Chapter 7: Vascular Modulatory Functions of Macrophages. Ioanna Keklikoglou and Michele De Palma; Editors: Subhra K. Biswas and Alberto Mantovani. Springer.

### 1.6.5 Lymphangiogenesis

In the embryonic development lymphatic vessels are established after the cardiovascular system, when subpopulations of endothelial cells commit to the lymphatic lineage (159). Lymphatic capillaries are unidirectional and are lined with a thin single layer of lymphatic endothelial cells (LECs) anchored to the ECM (160). These LECs have a discontinuous basement membrane and lack the coverage of pericytes or smooth muscle cells that makes them highly permeable to interstitial fluids and proteins and facilitates immune cell transmigration (161, 162). Hence, the function of lymphatic vessels includes the regulation of tissue fluid homeostasis, collection of macromolecules from tissues back to the blood circulation as well as the transport of immune cells (162).

As angiogenesis, lymphangiogenesis in adult tissues only occurs during tissue repair, inflammation or tumorigenesis (163). Both VEGF-C and -D are involved in physiological

and pathological lymphangiogenesis (164, 165). Whereas the deletion of *Vegfd* in mice results in slight reduction of lymphatic vessels, deletion of *Vegfc* leads to embryonic lethality. Thus, VEGF-C is essential for initial sprouting, migration and survival of LECs (166, 167). Tumor-associated lymphatic vessel growth is positively correlated to the overexpression of VEGF-C and -D in tumors (168, 169). Tumor-associated lymphatic vessels occur mostly at the tumor margin but may also occur intratumoral (169). LECs isolated from normal tissue compared to the ones from fibrosarcoma displayed significant differences in the expression of 790 genes. Among those, genes implicated in endothelial junctions, matrix and vessel growth were upregulated, while matrix proteins like collagens and fibrillin were downregulated (170, 171).

Lymphangiogenesis is also mediated by several subpopulations including tumor cells, stromal cells, activated platelets and infiltrating macrophages (172, 173). TAMs do not only secrete the lymphangiogenic factors VEGF-C and -D, they also respond to the chemotactic properties of VEGF-C as they express its receptor VEGFR-3 (174). Thus, TAMs have been associated with lymphangiogenesis in several studies (172, 175). The accumulation of TAMs significantly correlates with lymph vessel density in several tumor tissues (176). In human breast cancer, elevated levels of VEGF-C expressing TAMs are not only associated with increased lymph vessel density but also with lymph node metastasis and lymph vessel invasion (177). Interestingly, mechanistic evidence has been provided for the involvement of TNF- $\alpha$  activated TAMs in lymphangiogenesis and lymphatic metastasis. These TAMs mediate the observed effects by coordinating VEGF-C/VEGFR-3 signaling (178).

### **1.6.6 Invasion and Metastasis**

Invasion and metastasis is a multistep process that includes a cascade of different events. TAMs are one of the major players as they contribute at various steps (179). In the early stage of tumor development, macrophages are found in areas of basement membrane breakdown and in advanced tumors at the invasive front (106, 180). Tumor cells exploit the matrix deposition and remodeling capacities of TAMs that produce MMPs and other proteolytic enzymes (181, 182). For instance, the proteolytic capacity of MMP-9 releases matrix-bound VEGF-A that contributes to angiogenesis and metastasis (183, 184). Moreover, it has been shown that the release of MMPs by TAMs promotes tumor cell motility and invasiveness (148, 185-187). Thus, TAM-derived enzymes help in the establishment of a premetastatic and metastatic niche (84).

Intravasation also requires close interaction between tumor cells and TAMs. The release of CSF1 by tumor cells stimulates migration of CSF1R expressing macrophages and their production of epidermal growth factor (EGF) in turn activates invasion of EGFR expressing tumor cells. Depletion of either one of the factors of this paracrine signaling loop abrogates migration of both cell types *in vivo* (125, 188). Interestingly, Wyckoff and colleagues could show that macrophages guide tumor cells to the blood vessels and thereby increase migration and invasion (189). Additionally, TAMs increase tumor cell intravasation by contributing to vascular abnormalization creating leaky blood vessels. Thus, macrophage re-education of M2-like macrophages to M1-like macrophages inhibits endothelial cell proliferation and normalizes the vessel walls, which in turn prevents tumor cell intravasation and metastasis (190). However, at the metastatic site tumor cells recruit macrophages to also induce vessel permeability via the upregulation of VEGF-A enabling them to extravasate (191). These macrophages were characterized as CCR2<sup>+</sup>VEGFR1<sup>+</sup>Ly6C<sup>-</sup>F4/80<sup>+</sup> macrophages (192).

Tumor metastasis to distant organs and lymph nodes is well established. Tumor cells can metastasize to one or more distant sites either sequentially or in parallel. (193). However, there are still contradictory studies about the role of lymph nodes in metastasis. On one hand, it is reported that lymph nodes promote further spreading and on the other hand studies point out that lymph nodes only serve as distant sites of tumor cell colonization without any further dissemination (194-197). Nevertheless, there is one common consensus about lymph node metastasis being a key prognostic marker for patient survival and outcome (198, 199).

In conclusion, TAMs play an important role in the process of tumor metastasis and high infiltration of TAMs is correlated with increased tumor cell dissemination, malignancy and tumor grade in many types of cancer (92, 145, 200).

## **1.7 THERAPEUTIC IMPLICATIONS**

TAMs have emerged as attractive targets in anti-cancer therapy because they are the predominant cell type in many different cancers, they contribute in various ways to cancer progression and are associated with therapy resistance (73, 201, 202). Importantly, compared to cancer cells they are genetically stable and thus less susceptible to acquire therapeutic resistance (203). So far, several studies have been conducted targeting TAMs from different angles. These include inhibition of macrophage recruitment, macrophage re-education toward an anti-tumoral phenotype and suppression of macrophage survival (85, 202).

For instance, inhibition of CCL2 or VEGFR-2 reduces the infiltration of macrophages and inhibits tumor growth and angiogenesis (204, 205). Another very attractive target to block the recruitment of TAMs to tumor sites is the CSF1/CSF1R pathway. Ries and colleagues showed that blocking the CSF1R with the monoclonal antibody RG7155 reduces the recruitment of CSF1R<sup>+</sup>CD163<sup>+</sup> macrophages to the tumor in patients suffering from diffuse-giant sarcoma improving their clinical outcome (206). Furthermore, inhibition of CSF1R reprograms microglia from a pro-tumoral phenotype to an anti-tumoral phenotype that hampers glioma development and increases survival (91). However, CSF1R inhibition as a monotherapy does not always succeed. In preclinical models of breast cancer there was only an effect in tumor growth suppression when CSF1R inhibition was used in combination with other drugs, such as paclitaxel (207, 208). This different outcome points out the important fact that the composition of anti-cancer therapy is dependent on the cancer type and location and that monotherapy is not sufficient enough as it only delays tumor growth. Therefore, the combination with irradiation, anti-angiogenic therapies, adoptive T cell transfer and immune checkpoint inhibitors represents attractive partners (209-212).

Since it is known that M1-like macrophages execute anti-tumoral activity, re-educating pro-tumoral M2-like macrophages to anti-tumoral M1-like macrophages illustrates another attractive strategy. In this respect, several studies have shown anti-cancer potential using this approach. Coscia and colleagues have shown that zoledronic acid skews macrophages from an M2-like to an M1-like phenotype and thereby inhibits spontaneous mammary carcinoma development (213). Moreover, histidine-rich glycoprotein induces the re-education of TAMs toward an M1-like phenotype resulting in vessel normalization, increased vessel functionality and hampering of tumor growth and metastasis (190).

Finally, suppression of TAM survival has also been shown to improve therapeutic outcomes. Cieslewicz and colleagues developed an M2-like macrophage specific pro-apoptotic peptide (M2pep) that selectively killed TAMs in tumor-bearing mice improving their survival rates (214).

All these different approaches display high potential of using TAMs as a target in anti-cancer therapy and underline that the microenvironment is an important player to consider in successful treatment.

## 2 AIMS OF THE THESIS

The overall aim of this thesis is to elucidate the diverse functions of tumor-associated macrophages and microglia in tumorigenesis and metastasis. Understanding the mechanisms of how macrophages and microglia contribute to tumor malignancies in different microenvironments helps to develop tools that specifically target distinct subpopulations.

The thesis consists of the following four studies:

**Study I:** Exploring the effect of SEMA3A on the accumulation of anti-tumoral macrophages and their anti-tumor activity in mouse and human breast cancer.

**Study II:** Elucidating the role of macrophage- and tumor-derived VEGF-C on tumor metastasis in mouse and human breast cancer.

**Study III:** Studying the effect of Cripto-1 vaccination on metastasis and cancer stem cells in mammary carcinoma.

**Study IV:** Examining the function of macrophages and microglia in PDGFB-driven mouse and human glioma.

## 3 RESULTS AND DISCUSSION

### 3.1 TUMOR MOUSE MODELS

TAMMs, major players in tumor progression and metastasis, can be of an anti- or pro-tumoral phenotype that can either restrict or fuel tumor development at various steps. In this thesis we have investigated the anti- and pro-tumoral properties but also functional differences of TAMMs in breast cancer and glioma.

In **study I-III** we used the 4T1 mammary tumor mouse model to investigate tumor-associated macrophages and tumor cells in breast cancer evolvement and dissemination. 4T1 tumor cells were originally isolated from a single spontaneous tumor that arose in a BALB/cfC3H mouse and resembles human TNBC (215, 216). We engineered the cell line according to our aims to either overexpress SEMA3A, VEGF-C or cripto-1 and injected the cells orthotopically into the mammary fat pad of mice.

There are several advantages of using 4T1 tumor cells to study breast cancer. First, it is an easy transplantable cell line that is highly tumorigenic and invasive. As it is injected into the mammary fat pad it grows in the anatomically correct site. Second, 4T1 tumor cells spontaneously metastasize from the primary tumor mainly via the hematogeneous route to lung, liver, brain, bone and blood and to a lesser extent to the lymph nodes. Thus, 4T1 tumor cell dissemination is similar to that of human breast cancer. Finally, 4T1 tumor cells are resistant to 6-thioguanine enabling the quantification of metastasis in the mentioned distant organs (217). Of note, this model displays high accumulation of granulocytes into the tumors and lungs, which is a dissimilarity to human TNBC.

In **Study IV** we expanded our knowledge about macrophages by elucidating the functional role of microglia and their difference to macrophages in glioma development. We used the RCAS/*tv-a* mouse glioma model. Mice of this model harbor two genetic aberrations including the systemic depletion of the tumor suppressor Arf and the expression of the *tv-a* gene in glial progenitor cells that contain the nestin promoter. The *tv-a* gene encodes the receptor for the avian leukosis virus (ALV) and is normally only expressed in avian cells. The viral vector RCAS that is derived from the same virus is packaged with *pdgfb* and newborn mice are transduced with this virus allowing the selective transduction of glial progenitor cells that harbor the nestin promoter. Thus, those mice develop PDGFB driven N/*tv-a*;Arf<sup>-/-</sup> gliomas at high rates up to 100% incidence at 12 weeks of age (218, 219). This glioma model

is a well established and widely used model to study the mechanisms of glioma development and provides a life-like model of human glioma of grade II to IV (219).

## **3.2 STUDY I**

### **Guidance molecule SEMA3A restricts tumor growth by differentially regulating the proliferation of tumor-associated macrophages**

Semaphorin (SEMA)3A is a secreted protein that was originally discovered as an axon guidance regulator in establishing the neuronal network in the embryonic nervous system (220, 221). This protein is found in most human tissues and binds to its co-receptor neuropilin (NP)1 that forms receptor complexes with receptors of the Plexin family (222). However, in the last decade SEMA3A has been implicated in cancer development. Except for pancreatic cancer, where SEMA3A expression has been positively correlated to increased malignancy grade, this protein is down-regulated in several human cancers such as breast, gastric, epithelial ovarian and non-small cell lung cancer (119, 223-226).

Animal studies investigating the effect of SEMA3A on macrophages in the tumor microenvironment show contradictory results. Carrer and colleagues demonstrated that SEMA3A promotes vessel maturation through the recruitment of circulating NP1 expressing monocytes. These CD11b<sup>+</sup>NP1<sup>+</sup>Ly6G<sup>-</sup> monocytes, isolated either from the bone marrow or from SEMA3A-expressing muscles and directly injected into growing tumors, exerted anti-tumoral activity by induction of vessel normalization and inhibition of tumor growth (227). On the other hand, Casazza and colleagues reported that SEMA3A is induced in hypoxic areas and acts as an attractant for NP1-expressing TAMs. Once those TAMs have arrived in the hypoxic environment, they are entrapped inside the hypoxic niche and exert pro-angiogenic and immune-suppressive functions promoting tumor growth and metastasis (228).

However, knowing that SEMA3A expression is down-regulated in many different human cancer types, we also checked SEMA3A expression levels in different grades of human breast cancer samples. In concordance, we found a down-regulation of SEMA3A with disease progression. To further study the effect of SEMA3A on tumor development, we used the 4T1 tumor mouse model with 4T1 cells that overexpressed SEMA3A. Tumors overexpressing SEMA3A showed reduced tumor growth and burden. When we dissected the tumors and looked at the immune cell composition, we discovered an increased accumulation



of anti-tumoral M1-like macrophages and cytotoxic lymphocytes including T lymphocytes and NK cells.


The M1- and M2-like phenotypes of macrophages were determined by the expression of different cell surface markers using flow cytometry and gene expression profiles via RT-qPCR. Using flow cytometry we classified macrophages as CD11b<sup>+</sup>Ly6G<sup>-</sup> cells. On the basis of MHC class II expression we further subdivided macrophages into Ly6C<sup>lo</sup> MHC class II<sup>hi</sup> M1-like macrophages and Ly6C<sup>lo</sup> MHC class II<sup>lo</sup> M2-like macrophages. Additionally, we further verified our observation by investigating M1- and M2-markers that are involved in costimulation, antigen presentation and scavenging including CD11c, CD80, CD86, MHC class I and MRC1 in CD11b<sup>+</sup>F4/80<sup>+</sup> macrophages. Sorted CD11b<sup>+</sup>F4/80<sup>+</sup> macrophages from SEMA3A and CTR tumors were subjected to RT-qPCR looking at a range of cytokines and growth factors confirming our result of increased M1-like macrophage accumulation in SEMA3A tumors.

Based on our results and the fact that tumors are infiltrated by monocytes, we speculated that monocyte recruitment might account for the observed effect. However, there was no increase in monocyte frequency in the blood or tumor in SEMA3A overexpressing tumors. Furthermore, direct administration of SEMA3A to BMDMs did not affect their phenotype. Interestingly, publications about local macrophage proliferation led us to investigate if SEMA3A affected proliferation rather than recruitment of M1- and M2-like macrophages. Indeed, we could show both *ex vivo* and *in vivo* that SEMA3A mediates the expansion of M1-like macrophages, while it reduces the proliferation of M2-like macrophages. This effect was NP1 dependent and regulated Akt and MAPK signaling. Intriguingly, knockdown of NP1 in BMDMs mimicked the effect of SEMA3A on Akt and MAPK phosphorylation mediated by CSF1 stimulation. Thus, CSF1 stimulated M1-BMDMs showed increased Akt and MAPK phosphorylation after SEMA3A treatment or NP1 inhibition, while M2-BMDMs displayed reduced Akt and MAPK phosphorylation. NP1 can undergo signaling complexes with various members of the Plexin A family and therefore, we speculate that differential expression of Plexin A receptors and diverse dimerization complexes with NP1 could induce the opposing outcome in M1- and M2-macrophage proliferation. Further investigations are needed to explore the detailed mechanism of SEMA3A-mediated macrophage proliferation.

However, SEMA3A overexpressing tumors did not only show accumulation of anti-tumoral macrophages but also of cytotoxic CD8<sup>+</sup> T lymphocytes and NK cells. The increased recruitment and activation of CD8<sup>+</sup> T lymphocytes and NK cells was dependent on and mediated by SEMA3A via NP1 expression on macrophages. Nevertheless, the observed

reduced tumor growth of SEMA3A tumors was attributed to both macrophages and cytotoxic cells. Accordingly, upon depletion of the different populations SEMA3A lost its effect on tumor growth inhibition.

In sum, we show that SEMA3A mediates the proliferation of M1-like macrophages that creates a more pro-inflammatory microenvironment enhancing the recruitment and activation of cytotoxic CD8<sup>+</sup> T lymphocytes and NK cells and thereby inhibiting breast cancer progression (**Figure 9**).




**Figure 9.** SEMA3A enhances the expansion of M1-like macrophages and restricts the proliferation of M2-like macrophages. The selectively regulated proliferation recruits and activates cytotoxic CD8<sup>+</sup> T lymphocytes and NK cells and thereby inhibits tumor progression.

### 3.3 STUDY II

#### Macrophage-derived Vascular Endothelial Growth Factor-C decreases hematogeneous metastatic dissemination by normalizing the tumor vasculature

VEGF-C is a secreted dimeric glycoprotein and one of five family members in mammals (229). In adults VEGF-C expression is found in many different organs including heart, lung, pancreas and kidneys (230, 231). VEGF-C can bind to its two receptors VEGFR-2 that interacts with the co-receptor NP1 and VEGFR-3 that binds NP2 (230, 232, 233). The affinity of VEGF-C to either one of the receptors depends on the proteolytic cleavage. While non-processed VEGF-C activates VEGFR-3 signaling, mature VEGF-C binds with high

affinity to both VEGFR-2 and -3 and can induce the formation and activation of receptor heterodimers (**Figure 10**). These heterodimers have been reported to be present on lymphatic vessels but also on angiogenic sprouts pointing out a bimodal function for VEGF-C and its receptors (234-236). In fact, a study conducted by Tammela and colleagues showed that Tie2<sup>+</sup>VEGF-C<sup>+</sup>-macrophages accumulate behind tip cells at vascular branching points inducing endothelial cell differentiation via VEGFR-3 and Notch signaling. Thus, this finding indicates a function of VEGF-C-expressing macrophages in vessel maturation (237). Of note, processing of VEGF-C also affects its binding to the two co-receptors NP1 and NP2. While the partially processed protein binds to both NP1 and NP2, only the fully mature protein can bind NP2 showing another layer of complexity in VEGF signaling (238).



**Figure 10. VEGF-C binding specificities and VEGFR signaling complexes.** Adapted and modified from (239).

However, VEGF-C expression has been detected in several different types of human tumors including breast, cervix, colon, lung, prostate and stomach (240-242). Although VEGF-C expression is predominantly correlated to lymphangiogenesis, it has been shown that tumor cell-secreted VEGF-C binds to VEGFR-3 that is upregulated on angiogenic blood vessels in breast cancer (243-245). Several different types of tumors express VEGF-C but TAMs have been reported to be a major source of this protein and as mentioned before they correlate to lymphatic microvessel density, lymphangiogenesis, invasion and lymph node metastasis in several types of cancer (172, 174, 177, 246, 247).


As tumor dissemination occurs later during cancer progression and macrophages are active participants in this process, it would be natural to assume that macrophages and tumor cells secrete increased levels of VEGF-C leading to lymph node metastasis in advanced breast cancer. However, when we looked at VEGF-C expression in ER<sup>+</sup> grade I and III ductal carcinoma and grade III TNBC, we found unexpectedly decreased VEGF-C expression by CD68<sup>+</sup> macrophages in grade III compared to grade I breast cancer. Moreover, the overall expression of VEGF-C within the tumor seemed to be similar throughout the different grades. This evidence prompted us to further investigate if tumor- and macrophage-derived VEGF-C might be differentially regulated during tumor progression and thereby influence the route of tumor cell dissemination.

To investigate our hypothesis we chose to use the 4T1 tumor mouse model. As mentioned earlier this mouse model resembles TNBC and mainly metastasizes via the hematogeneous and not lymphangiogenic route. Accordingly, we expected high levels of VEGF-A in macrophages and tumor cells. Therefore, macrophages and tumor cells were sorted from 4T1 tumors and VEGF-A, -C and -D expression levels were analyzed by RT-qPCR. Intriguingly, macrophages only expressed VEGF-A while VEGF-C and -D were undetected. However, tumor cells expressed both VEGF-A and VEGF-C to a similar extent. Nevertheless, it was evident that macrophages expressed 10 times higher levels of VEGF-A compared to tumor cells, so that we speculated that macrophage-derived VEGF-A rather than tumor-derived VEGF-A dictates the preferred hematogeneous route for 4T1 cancer cell dissemination. We further hypothesized that if we engineer macrophages to overexpress VEGF-C, we might be able to change the preferable route of dissemination from blood vessels to lymphatic vessels.

To test our theory, we engineered chimeric mice that had been transplanted with bone marrow-derived stem cells to express VEGF-C using lentiviral-mediated gene transduction. These chimeric mice were then inoculated with 4T1 tumor cells. In comparison, to test what effect tumor-derived VEGF-C has, 4T1 tumor cells were also engineered to overexpress VEGF-C by lentiviral-mediated gene transduction and injected into the fat pad of naïve mice. Neither macrophage- nor tumor-derived VEGF-C altered tumor growth or burden compared to their respective controls. However, tumor- and macrophage-derived VEGF-C had a different effect on lung metastasis. While both tumor- and macrophage-derived VEGF-C increased lymph node metastasis, only macrophage-derived but not tumor-derived VEGF-C decreased lung metastasis. In fact, tumor-derived VEGF-C increased pulmonary metastasis but not as a consequence of increased angiogenesis. Consequently, we considered another important factor contributing to tumor spreading, namely vessel dysfunctionality. Therefore,

we determined blood vessel perfusion, pericyte coverage and hypoxia. Interestingly, macrophage-derived VEGF-C increased vessel perfusion, pericyte coverage and decreased hypoxia while tumor-derived VEGF-C had the opposite effect. Improved vessel functionality is attributed to the M1-like phenotype but VEGF-C expression in macrophages or tumor cells did not shift the macrophage phenotype toward an M1- or M2-like phenotype. Moreover, neither the infiltration of cytotoxic CD8<sup>+</sup> T lymphocytes nor NK cells was altered. Further investigations are necessary to elucidate the exact mechanism of improved vessel functionality by macrophage-derived VEGF-C. It would be interesting to elaborate if processing of VEGF-C and the binding to its two receptors VEGFR-2 and -3 is regulated differentially by tumor cells and macrophages leading to the various effects on vessel functionality.

Taking all results in perspective, we show that macrophage-derived VEGF-C normalizes the tumor vasculature and thereby decreases pulmonary metastasis while tumor-derived VEGF-C increases vessel abnormalization and lung metastasis (**Figure 11**).



**Figure 11.** Macrophage-derived VEGF-C normalizes the blood vessels and thereby decreases lung metastasis, while tumor-derived VEGF-C decreases vessel functionality leading to increased pulmonary metastasis.

### 3.4 STUDY III

#### **Cripto-1 plasmid DNA vaccination targets metastasis and cancer stem cells in murine mammary carcinoma**

Cripto-1 is a protein that plays an important role during embryonic development contributing to the undifferentiated status of human and mouse embryonic stem cells (248). However, more than 50% of human cancers, including breast cancer, show elevated Cripto-1 expression levels and its presence on cancer cells has been associated with the cancer stem cell (CSC) phenotype (249, 250). CSCs are a subpopulation that initiates tumor growth, shows enhanced metastatic potential, is resistant to radiation and chemotherapy and responsible for relapse (249, 251). Indeed, cripto-1 has been shown to be involved in proliferation, angiogenesis and invasion and high levels are associated with colorectal metastasis and poor prognosis in colon and breast cancer (252-256). Therefore, Cripto-1 displays an interesting candidate to specifically target metastatic cancer cells. For instance, in a murine melanoma model Cripto-1 vaccination elicited a specific cytotoxic CD8<sup>+</sup> T lymphocyte mediated protective response by reducing primary tumor burden and lung metastasis (257).


Hence, we wanted to investigate if Cripto-1 vaccination also had a protective effect in mammary breast cancer progression and metastasis. Therefore, different cell lines were checked for Cripto-1 expression. Cripto-1 could be detected in the highly metastatic 4T1 tumor cell line and in TUBO cells that show a cancer stem cell like phenotype. However, Cripto-1 expression in 4T1 cells was very low and not detectable via flow cytometry, so that we decided to generate a stable Cripto-1 expressing 4T1 cell line that could be used in our *in vivo* studies. Before inoculation of these cells into the mammary fat pad, mice were vaccinated with Cripto-1 encoding DNA plasmids. Intriguingly, Cripto-1 vaccination did not only reduce tumor growth and burden but also decreased lung metastasis.

We then wondered how this effect was mediated. Vaccination goes along with cell- and antibody-mediated immunity. Accordingly, we speculated that Cripto-1 vaccination could evoke a humoral response in mice. Indeed, we detected Cripto-1 specific antibodies in serum of vaccinated mice that could bind to Cripto-1 expressing 4T1 cells. Moreover, most of the antibodies belonged to the subclasses IgG2a and IgG2b that in mice mediate antibody-dependent cell-mediated cytotoxicity (ADCC) by NK cells. To test if Cripto-1 specific antibodies induce ADCC, we investigated if serum from Cripto-1 vaccinated mice could increase NK cell cytotoxicity. In fact, pre-activated NK cells showed increased lysis of

Cripto-1 expressing 4T1 cells in the presence of serum from Cripto-1 vaccinated mice *in vitro*.

Furthermore, to verify our observations of reduced tumor and metastatic burden in our validation model (Cripto-1 expressing 4T1 tumor mouse model), we used the more clinical relevant BALB-neuT mouse model. Intriguingly, despite Cripto-1 negative primary tumors, we observed a protective effect of Cripto-1 vaccination on the metastatic burden in the lungs. Furthermore, Cripto-1 vaccination effectively decreased tumor growth of TUBO cells that show CSC characteristics and even led to tumor clearance, as three out of eleven mice stayed tumor free for more than 60 days.

Taking all together, Cripto-1 vaccination induced a humoral response facilitating NK cell-mediated ADCC and the reduction of lung metastasis (**Figure 12**).



**Figure 12.** Cripto-1 DNA plasmid vaccination decreases the metastatic burden in lungs of mice harboring Cripto-1 positive and negative primary tumors. NK cells show enhanced killing of Cripto-1 expressing 4T1 tumor cells in the presence of Cripto-1 serum.

### 3.5 STUDY IV

#### **Microglia induce PDGFRB expression in glioma cells to enhance their migratory capacity**

Platelet-derived growth factor (PDGF)B and one of its receptors, PDGFRB, are essential for physiological processes like proliferation, migration and blood vessel development but have also been linked to pathological conditions, such as glioma development (258-260). This signaling pathway is mostly linked to tumor-associated vasculature and angiogenesis. For instance, PDGFB expression by endothelial cells regulates vessel permeability and vascular maturation in extracranial solid tumors (260, 261). Moreover, tumor cells that overexpress PDGFB induce displacement of PDGFRB<sup>+</sup> pericytes from endothelial cells, which increases vessel dysfunctionality and metastatic dissemination (262). However, even though not one of the most prevalent alterations, aberrations in PDGFB expression and/or PDGFRB phosphorylation are observed in gliomas and GBMs (218, 260, 263). Interestingly, there is one study reporting that PDGFRB is commonly expressed in cultured patient-derived GBM cells, especially by self-renewing GBM stem cells (264).

At the time when this study was initiated PDGFB-driven glioma was not well characterized with regards to the tumor and microenvironmental signature. Using the RCAS/*tv-a* mouse glioma model we defined PDGFB driven N/*tv-a*;Arf<sup>-/-</sup> gliomas and elucidated the functional properties of microglia and macrophages.


In collaboration with a neuropathologist, according to histopathological characteristics of human gliomas, PDGFB driven N/*tv-a*;Arf<sup>-/-</sup> mouse gliomas were divided into grade II- to grade IV-like gliomas, where grade IV-like gliomas showed features of human GBMs. Moreover, these low- and high-grade gliomas could be identified as astrocytomas based on their expression of GFAP.

We then continued to elaborate on vessel functionality and analyzed vessel perfusion and hypoxia. With increased grade of malignancy the number of perfused vessels was significantly decreased while hypoxia was markedly increased. Since pericyte coverage of blood vessels is important for vessel maturation and function, we assumed that high-grade gliomas have decreased pericyte coverage and immunostained the gliomas with the pericyte marker  $\alpha$ -smooth muscle actin ( $\alpha$ -SMA). Surprisingly,  $\alpha$ -SMA<sup>+</sup> cells accumulated with increased malignancy and these cells were rarely found in conjunction with vessels but instead spread throughout the tumors. This finding prompted us to further investigate if those  $\alpha$ -SMA<sup>+</sup> cells were actually pericytes. Therefore, we checked different mesenchymal/pericyte




and tumor cell markers and could identify those cells as a subpopulation of tumor cells that expressed  $\alpha$ -SMA and PDGFRB.

Considering that  $\alpha$ -SMA<sup>+</sup> and PDGFRB<sup>+</sup> glioma cells were scattered throughout the tumor mass, we wondered if these cells were associated to other types of stromal cells, such as TAMMs. Indeed, we found IBA1<sup>+</sup> TAMMs in close proximity to  $\alpha$ -SMA<sup>+</sup> and PDGFRB<sup>+</sup> glioma cells. We then continued investigating PDGFRB expression on glioma cells and their co-localization with TAMMs. In concordance, we found increased PDGFRB expression on glioma cells in high-grade gliomas compared to low-grade gliomas. Along with this finding we observed increased IBA1<sup>+</sup> TAMM accumulation that could be correlated to PDGFRB expression on glioma cells. Between 50-65% of those glioma cells were in direct physical contact to IBA1<sup>+</sup> TAMMs. However, we were curious about how relevant our observations in mouse astrocytomas would be for human astrocytomas. Hence, we looked at PDGFRB expression on glioma cells and TAMM accumulation in grade II to IV human astrocytomas. In fact, we could verify the correlative accumulation of TAMMs and increased PDGFRB expression on glioma cells throughout the grades of human astrocytomas (**Figure 13**).



**Figure 13.** Mouse and human astrocytomas show increased expression of PDGFRB on glioma cells with increased grade of malignancy, which correlates to the accumulation of TAMMs.

Based on these results, we hypothesized that TAMMs could induce PDGFRB expression on glioma cells. As mentioned earlier, under physiological conditions, microglia are the abundant cell type in the brain. However, when pathological malignancies occur, the blood brain barrier is disrupted and monocytes can infiltrate the tumor. Therefore, we performed co-culture experiments with glioma stem cells together with either microglia or BMDMs that were polarized to an M1-like phenotype (LPS+IFN $\gamma$ ) or M2-like phenotype (IL-4+TGF- $\beta$ +IL-10) *ex vivo*. To our surprise, only cell-to-cell contact with microglia but not BMDMs could induce glioma cells to express PDGFRB. Moreover, even though both M1- and M2-like microglia could induce PDGFRB expression, M2-like microglia induced significantly higher expression levels of this receptor. Importantly, only M2-like microglia were able to increase the migratory capacity of glioma cells via the upregulation of PDGFRB. Of note, BMDMs that have been pre-conditioned with microglia-derived factors also failed to induce PDGFRB expression on glioma stem cells. In conclusion, microglia but not BMDMs induce PDGFRB expression on glioma cells and thereby enhance their migratory capacity (**Figure 14**).



**Figure 14.** Microglia induce PDGFRB expression on glioma cells and thereby increase their migratory capacity. Macrophages fail to induce PDGFRB expression on glioma cells.

The finding that BMDMs failed to induce PDGFRB expression on glioma cells *in vitro*, sheds more light on the functional differences between microglia and macrophages that have not been fully discovered yet. This is especially important in therapy treatment because approaching these two subpopulations differentially might present new possibilities in glioma treatment.

## 4 CONCLUSIONS

Cancer is a complex disease that involves the immune system at every step of its growth and spread to secondary organs. The wide variety of mutations and differential microenvironmental signatures that may be encountered throughout the various cancer types make it very difficult to successfully treat the disease. Macrophages and microglia are abundant cells in all types of cancers. Whether these cells are allies or enemies to cancer depends on their microenvironmental surroundings and stimuli. In this thesis we have demonstrated the diverse functions of macrophages and microglia in cancer evolvment and invasion, and have shown that modulating, rather than depleting distinct subpopulations of macrophages and microglia, offers more opportunities to develop successful strategies in anti-cancer therapies. Moreover, we have illustrated that there are functional dissimilarities between macrophages and microglia. These dissimilarities provide evidence that these two populations not only differ ontologically but also behave differentially when interacting with tumor cells. These findings may facilitate better understanding of the efficiency and outcome of therapy treatments.



## 5 ACKNOWLEDGEMENTS

I am very grateful for the opportunity to perform my PhD at Karolinska Institute. At the beginning of my doctoral studies I was told that the years of being a PhD student resemble a roller coaster with many up and downs. It truly was a roller coaster for me but at the end of my ride I can honestly say that I am happy about my decision to do it and to meet so many people that supported me throughout my time at CCK.

First of all I would like to thank my main supervisor **Charlotte**. I very much appreciate the opportunity you gave me to perform my PhD in your lab. Thank you for all your support and availability during all those years. I learned so much during this time and you have helped me develop and getting stronger not only on a scientific but also personal level.

My co-supervisors **Andreas, Arne, Ola, Tatiana** and mentor **Marianne**, thank you for your guidance. Especially **Andreas**, thank you for being there for me when I needed it and cheering me up. I am very grateful for your support.

I would like to thank my examiners **Myriam Aouadi, Lars-Gunnar Larsson** and **Lasse Jensen** and my opponent **Claire Lewis** for being part of my examination committee.

My wonderful lab members that created the best atmosphere at work. **Majken**, it has been such a pleasure to meet you and become friends with you. Since my first day in the group you were always there for me and you never let me down. Without you I would never have experienced such an amazing time. I am very happy that we became friends and that you included me into your family. I enjoyed every moment. I also could not imagine a better colleague. **Margarita**, even though you joined when I was almost halfway through, I enjoyed very much to get another wonderful colleague and friend. I could always count on you and you were there when I needed you. Sharing good and difficult times and becoming friends beyond work gave me great support. I enjoyed spending time with you and I appreciate this very much.

**Jeanette** and **Emma**, thank you for the nice time, help and all your valuable contributions. It was a pleasure and fun time working with you. **Pradeepa**, your smile and happy attitude were always contagious. Working in the lab with you was like having a smiling sun next to me. **Dennis** and **Laura**, both of you always wished a good morning with a smile on your face. Your positive and motivated attitude spread great warmth. I had so much fun with you and I was impressed how dedicated you worked in the lab. Thank you for all your contributions and staying friends beyond lab work. **Hui** and **Stina**, even though we spent

little time together it was very nice having you as colleagues and get to know another two very nice girls.

All the present and past floor members on the first floor created a stimulating atmosphere and fun fikas.

**Andreas Lundqvist group:** **Kristina**, thank you for being a friend and coming by to the lab or office even though it was just to ask how I am doing and to give me a hug. Talking to you made me always feel better. I could always count on you when I needed help. I appreciate this very much. Thanks to **Dhifaf, Erik, Chen, Neo, Ying** and **Veronika**.

**Rolf Kiessling group:** Many thanks for the good times to **Rolf, Yago, Ulrika, Stina, Jeroen, Takahiro, Marteen, Mao, Yuya** and **Tanja**.

**Ola Larsson group:** **Laia**, we did not only become office and cell lab mates but also roommates and friends. Thank you for inviting me to your hometown, organizing my surprise birthday dinner and letting me be a part of Èlias life. I am very grateful for all your support. **Vincent, Shuo, Julie, Johannes, Christian** (I use the small pan almost every day ☺) and **Baila** thank you for the fun times.

Thank you for a friendly atmosphere and nice talks to **Dalianis** and **Mellstedt groups:** **Tina, Linnéa, Cinzia, Torbjörn, Anders, Shahrzad, Nikos, Andreas, Cecilia, Nathalie, Kia, Barbro, Ann, Amir, Mohammad, Fariba** and **Amineh**. **Li-Sophie**, it was very nice chatting with you on our way to CCK. **Tom**, you are always a sincere and friendly person. I appreciate our fun and relaxing conversations.

**Pär Nordlund group:** **Smaranda, Sue-Li** and **Susanne**, thank you for having so many great talks and discussions when sharing office. **Henriette** and **Anderson**, it was nice having our short conversations about running and soccer. Thanks to **Pär, Olga, Sara** and **Anette**.

**Elle, Elisabeth, Sören** and **Eva-Lena**, you are the soul of CCK. Only with your help and contribution it is such a pleasure to work here!

I also want to show my gratitude to **Inger, Anna, Susanne, Juan** and **KI-fixit** for helping out with projects and problems.

Thanks to all my CCK and KI colleagues and collaborators for giving me happy and enjoyable moments.

**Xingmei**, thank you for all your optimism and making me believe in myself. It was very nice to work with you. **Bob**, thank you for listening and a fruitful collaboration.

**John, Anne-Laure and Christina**, you were always flexible and accommodating during the sorting times. I am very grateful for such a smooth collaboration. **Anne-Laure and Christina**, thank you for all the fun talks. I enjoyed them very much. **Lina**, I want to give you many thanks for always answering my questions and giving me very valuable suggestions. **Ioannis**, you are a very enthusiastic person that shows great interest and excitement about research. It is difficult not to share this when talking and working with you. **Susi**, cooking slides made so much more fun with having you there for a small chat. **Xuan and Johanna** working with you side by side was more working with friends instead of collaborators. Thank you for the great interactive work. **Nikos and Dimitris**, with you in the cell lab it never became boring. I had such a great and fun time working and sharing all kinds of gossip with you. **Sophia**, our coincidental encounters in the staircase or on the floors very often ended up in long conversations that I enjoyed very much. Sharing our experiences helped me a lot.

Thank you for nice collaborations to **Hrvoje Miletic, Per Enger, Margareta Wilhem, Rainer Heuchel** and all the interactive work and scientific exchange with **Arne Östman and Lars Holmgren groups. Carina, Alessandro, Pablo, Aravindh, Pedro, Evelyn, Yuan and Vicky**, I enjoyed all our scientific but also personal chats very much.

**Jonas Bergh and group**, I appreciate our collaborative work and meetings.

**Conan Hom**, thank you for taking your time and giving me great and valuable suggestions.

Ich möchte mich von ganzem Herzen bei meiner Familie und meinen Freunden aus Deutschland bedanken. Meine Erfahrung hier in Stockholm hat mir gezeigt, dass es nichts Schöneres auf der Welt gibt, als Zeit mit den Menschen zu verbringen, die mich so akzeptieren und schätzen wie ich bin, egal wo ich im Leben stehe. Zeit ist das kostbarste was man hat. Ich freue mich unglaublich darauf mehr Zeit mit euch zu verbringen. Die nachfolgenden Worte geben nur einen Bruchteil von der Dankbarkeit wieder, die ich für jeden Einzelnen von euch empfinde.

**Mama und Papa**, es gibt keine Worte, die beschreiben können wie dankbar ich euch bin für all das was ihr für mich getan habt. Die unerlässliche und unermüdliche Liebe und Unterstützung ist für euch immer selbstverständlich. Ihr lasst mich meinen Weg finden und gibt mir die Freiheit mich zu einem selbständigen und unabhängigen Menschen zu

entwickeln. In schwierigen Zeiten steht ihr mir ohne eine Sekunde zu zögern immer zur Seite. Ich kann immer auf eure Hilfe zählen, egal ob uns Länder oder Kontinente trennen. Ohne euch wäre ich niemals da wo ich jetzt bin. Ihr gibt mir Kraft, Mut und Zuversicht. Mit euch an meiner Seite habe ich das Gefühl, dass ich alles schaffen kann. Ihr bedeutet die Welt für mich und ich liebe euch über alles!

**Bruderherz**, ich könnte mir keinen besseren Bruder vorstellen als dich. Ich bewundere dich immer sehr für deine Hingabe und Hartnäckigkeit Probleme zu lösen. Wenn ich deine Hilfe brauche, nimmst du dir Zeit und ich kann immer auf dich zählen. Du nimmst mich so wie ich bin. Es ist schön zu wissen, dass du für mich da bist, auch wenn es nur darum geht eine Runde an der Rur zu laufen. Ich möchte dir von ganzem Herzen für alles danken.

**Schwesterherz**, die Distanz hat uns keinen Abbruch getan, sondern im Gegenteil noch enger zusammengeschweißt. Du hörst mir zu, bist da wenn ich dich brauche, verurteilst mich nicht und unterstützt mich in meinen Entscheidungen. Ich bewundere dich sehr für dein künstlerisches Talent, das mir schon so viel Freude bereitet hat. Du bist für mich ohne Frage die beste Schwester der Welt. Auch dir möchte ich von ganzem Herzen für alles danken.

**Housseem**, schon seit vielen Jahren gehörst du zum festen Bestandteil unserer Familie. Mit dir wird es nie langweilig und es gibt immer etwas zu lachen. Mit dir kann man immer eine unbeschwerte und entspannte Zeit genießen. Danke dafür.

**Cagla**, vor 15 Jahren haben wir uns getroffen und seit dem so viel gemeinsam erlebt. Jeder von uns beiden ist seinen Weg gegangen und doch haben wir uns nie verloren. Du bist immer für mich da, wenn ich dich brauche und du hast immer alles mit mir gemeinsam durchgestanden ohne mich jemals für etwas zu verurteilen. Du gehörst zu den Menschen, die immer an mich glauben. Dafür möchte ich dir von Herzen danken.

**Katja**, du gehörst ohne Frage zu meinem engsten Freundeskreis dazu. Wir sind uns nicht nur in vielerlei Hinsicht sehr ähnlich, sondern teilen auch sehr viele gemeinsame Ansichten. Du verstehst und unterstützt mich. Auch wenn wir uns in den letzten Jahren nicht häufig sehen konnten, ist es immer vertaut, wenn wir es schaffen uns in Stockholm oder Froitzheim zu treffen. Danke für all deine selbstlose Unterstützung, deinen Glauben an mich und dafür, dass du mich so schätzt wie ich bin.

**Dominique**, ab dem ersten Tag, an dem wir uns im Audimax begegnet sind, sind wir uns 3 Jahre nicht von der Seite gewichen und haben sehr viele lustige Momente gehabt und Diskussionen geführt. Du hast meine Studienzeit sehr bereichert und ich bin sehr dankbar



dafür, dass wir darüber hinaus so gute Freunde geblieben sind. Ich möchte dich nicht missen und schätze es sehr.

**Xenia**, wir kennen uns noch aus der Schulzeit und waren nicht nur ein “Dream Team” im Volleyball. Durch meine Leidenschaft habe ich dich kennengelernt und durch dich habe ich meine Leidenschaft wieder entdeckt. Wir haben sehr viel erlebt und uns letztendlich nicht aus dem Blick verloren. Es ist schön zu wissen, dass du immer für mich da bist und unsere Freundschaft so sehr schätzt. Das bedeutet mir sehr viel. Du und **Sven**, ihr empfangt mich immer mit offenen Armen und lasst es mir an nichts fehlen. Ich möchte mich für alles von Herzen bedanken.

**Mike**, uns verbindet eine sehr lange Vergangenheit, wo wir sehr viel geteilt, erlebt und gelernt haben. Wie das Leben so spielt, sollte es nicht bei dieser Vergangenheit bleiben. Was das Leben für uns noch bereit hält, weiß ich nicht, aber wir können es gemeinsam in der Gegenwart und Zukunft herausfinden. Danke für all deine Unterstützung, deine selbstlose Hilfe und deinen Glauben an mich. Du bringst mich immer zum Lachen und bist für mich da. Das alles bedeutet mir sehr viel.

**Desi**, ich bin unheimlich froh, dass wir uns wiedergefunden haben. Ich genieße es immer sehr mit dir, Dustin und Jayden Zeit zu verbringen und einfach mal ein spontanes Frühstück bei Claßen zu genießen. Danke für alles was du für mich getan hast und dafür, dass du immer für mich da bist. Mit dir kann ich Lachen und Weinen. Das bedeutet mir sehr viel.

**Olla, Micky, Julia, Bee, Valla, Johann und Markus**, jedes Mal wenn wir uns alle treffen, haben wir sehr viel Spaß. Es bleibt nie langweilig und es gibt immer sehr viel zu lachen. In unserer Runde können wir ausgelassen Blödsinn machen und den Stress drum herum vergessen. Danke für all diese ausgelassen Abende, die mich die Probleme immer für ein Weilchen vergessen lassen. Ich bin sehr froh euch alle zu haben.

**Ralf Gath**, während der Schulzeit begegnet jeder Schüler einem Lehrer, der ihn prägt und auf dem Weg „erwachsen zu werden“ interessiert begleitet. Du warst dieser Lehrer für mich. Ich bin dir sehr dankbar für all deine Unterstützung. Du hast nie etwas im Gegenzug gefordert. Die Volleyball AG war das Highlight meiner Schulzeit und immer das worauf ich mich am meisten gefreut habe.

**Martin**, wir kennen uns jetzt schon ein paar Jahre, in denen unsere Freundschaft stetig gewachsen ist. Es ist immer sehr nett mit dir über Gott und die Welt zu sprechen. Ich danke dir für die aktive Interesse an meinem Leben.

## 6 REFERENCES

1. Hanahan D, Weinberg RA. The hallmarks of cancer. *Cell*. 2000;100(1):57-70.
2. Hanahan D, Weinberg RA. Hallmarks of cancer: the next generation. *Cell*. 2011;144(5):646-74.
3. Hutchinson L. Breast cancer: challenges, controversies, breakthroughs. *Nat Rev Clin Oncol*. 2010;7(12):669-70.
4. Becker S. A historic and scientific review of breast cancer: The next global healthcare challenge. *Int J Gynaecol Obstet*. 2015;131 Suppl 1:S36-9.
5. Polyak K. Heterogeneity in breast cancer. *J Clin Invest*. 2011;121(10):3786-8.
6. Rakha EA, Reis-Filho JS, Baehner F, Dabbs DJ, Decker T, Eusebi V, et al. Breast cancer prognostic classification in the molecular era: the role of histological grade. *Breast Cancer Res*. 2010;12(4):207.
7. Giuliano AE, Edge SB, Hortobagyi GN. Eighth Edition of the AJCC Cancer Staging Manual: Breast Cancer. *Ann Surg Oncol*. 2018;25(7):1783-5.
8. Hon JD, Singh B, Sahin A, Du G, Wang J, Wang VY, et al. Breast cancer molecular subtypes: from TNBC to QNBC. *Am J Cancer Res*. 2016;6(9):1864-72.
9. Prat A, Adamo B, Cheang MC, Anders CK, Carey LA, Perou CM. Molecular characterization of basal-like and non-basal-like triple-negative breast cancer. *Oncologist*. 2013;18(2):123-33.
10. Rakha EA, Reis-Filho JS, Ellis IO. Basal-like breast cancer: a critical review. *J Clin Oncol*. 2008;26(15):2568-81.
11. Voduc KD, Cheang MC, Tyldesley S, Gelmon K, Nielsen TO, Kennecke H. Breast cancer subtypes and the risk of local and regional relapse. *J Clin Oncol*. 2010;28(10):1684-91.
12. Liedtke C, Mazouni C, Hess KR, Andre F, Tordai A, Mejia JA, et al. Response to neoadjuvant therapy and long-term survival in patients with triple-negative breast cancer. *J Clin Oncol*. 2008;26(8):1275-81.
13. Anders CK, Carey LA. Biology, metastatic patterns, and treatment of patients with triple-negative breast cancer. *Clin Breast Cancer*. 2009;9 Suppl 2:S73-81.
14. Tseng LM, Hsu NC, Chen SC, Lu YS, Lin CH, Chang DY, et al. Distant metastasis in triple-negative breast cancer. *Neoplasma*. 2013;60(3):290-4.
15. He M, Zhang JX, Jiang YZ, Chen YL, Yang HY, Tang LC, et al. The lymph node ratio as an independent prognostic factor for node-positive triple-negative breast cancer. *Oncotarget*. 2017;8(27):44870-80.
16. Liu N, Yang Z, Liu X, Niu Y. Lymph node status in different molecular subtype of breast cancer: triple negative tumours are more likely lymph node negative. *Oncotarget*. 2017;8(33):55534-43.
17. Schwartzbaum JA, Fisher JL, Aldape KD, Wrensch M. Epidemiology and molecular pathology of glioma. *Nat Clin Pract Neurol*. 2006;2(9):494-503; quiz 1 p following 16.
18. Agnihotri S, Burrell KE, Wolf A, Jalali S, Hawkins C, Rutka JT, et al. Glioblastoma, a brief review of history, molecular genetics, animal models and novel therapeutic strategies. *Arch Immunol Ther Exp (Warsz)*. 2013;61(1):25-41.
19. Hoshida R, Jandial R. 2016 World Health Organization Classification of Central Nervous System Tumors: An Era of Molecular Biology. *World Neurosurg*. 2016;94:561-2.
20. Louis DN, Perry A, Reifenberger G, von Deimling A, Figarella-Branger D, Cavenee WK, et al. The 2016 World Health Organization Classification of Tumors of the Central Nervous System: a summary. *Acta Neuropathol*. 2016;131(6):803-20.
21. Lin N, Yan W, Gao K, Wang Y, Zhang J, You Y. Prevalence and clinicopathologic characteristics of the molecular subtypes in malignant glioma: a multi-institutional analysis of 941 cases. *PLoS One*. 2014;9(4):e94871.
22. Cancer Genome Atlas Research N. Comprehensive genomic characterization defines human glioblastoma genes and core pathways. *Nature*. 2008;455(7216):1061-8.

23. Hambardzumyan D, Bergers G. Glioblastoma: Defining Tumor Niches. *Trends Cancer*. 2015;1(4):252-65.
24. Reifemberger G, Wirsching HG, Knobbe-Thomsen CB, Weller M. Advances in the molecular genetics of gliomas - implications for classification and therapy. *Nat Rev Clin Oncol*. 2017;14(7):434-52.
25. Yatim KM, Lakkis FG. A brief journey through the immune system. *Clin J Am Soc Nephrol*. 2015;10(7):1274-81.
26. Iwasaki A, Medzhitov R. Control of adaptive immunity by the innate immune system. *Nat Immunol*. 2015;16(4):343-53.
27. McComb S, Thiriot A, Krishnan L, Stark F. Introduction to the immune system. *Methods Mol Biol*. 2013;1061:1-20.
28. Perry VH, Hume DA, Gordon S. Immunohistochemical localization of macrophages and microglia in the adult and developing mouse brain. *Neuroscience*. 1985;15(2):313-26.
29. Gordon S, Taylor PR. Monocyte and macrophage heterogeneity. *Nat Rev Immunol*. 2005;5(12):953-64.
30. Martinez FO, Sica A, Mantovani A, Locati M. Macrophage activation and polarization. *Front Biosci*. 2008;13:453-61.
31. Yin J, Valin KL, Dixon ML, Leavenworth JW. The Role of Microglia and Macrophages in CNS Homeostasis, Autoimmunity, and Cancer. *J Immunol Res*. 2017;2017:5150678.
32. London A, Cohen M, Schwartz M. Microglia and monocyte-derived macrophages: functionally distinct populations that act in concert in CNS plasticity and repair. *Front Cell Neurosci*. 2013;7:34.
33. Aarum J, Sandberg K, Haerberlein SL, Persson MA. Migration and differentiation of neural precursor cells can be directed by microglia. *Proc Natl Acad Sci U S A*. 2003;100(26):15983-8.
34. Walton NM, Sutter BM, Laywell ED, Levkoff LH, Kearns SM, Marshall GP, 2nd, et al. Microglia instruct subventricular zone neurogenesis. *Glia*. 2006;54(8):815-25.
35. Wake H, Moorhouse AJ, Jinno S, Kohsaka S, Nabekura J. Resting microglia directly monitor the functional state of synapses in vivo and determine the fate of ischemic terminals. *J Neurosci*. 2009;29(13):3974-80.
36. Wynn TA, Chawla A, Pollard JW. Macrophage biology in development, homeostasis and disease. *Nature*. 2013;496(7446):445-55.
37. Schulz C, Gomez Perdiguero E, Chorro L, Szabo-Rogers H, Cagnard N, Kierdorf K, et al. A lineage of myeloid cells independent of Myb and hematopoietic stem cells. *Science*. 2012;336(6077):86-90.
38. Zhao Y, Zou W, Du J, Zhao Y. The origins and homeostasis of monocytes and tissue-resident macrophages in physiological situation. *J Cell Physiol*. 2018;233(10):6425-39.
39. Kierdorf K, Erny D, Goldmann T, Sander V, Schulz C, Perdiguero EG, et al. Microglia emerge from erythromyeloid precursors via Pu.1- and Irf8-dependent pathways. *Nat Neurosci*. 2013;16(3):273-80.
40. Gautier EL, Yvan-Charvet L. Understanding macrophage diversity at the ontogenic and transcriptomic levels. *Immunol Rev*. 2014;262(1):85-95.
41. Gentek R, Molawi K, Sieweke MH. Tissue macrophage identity and self-renewal. *Immunol Rev*. 2014;262(1):56-73.
42. Bain CC, Bravo-Blas A, Scott CL, Perdiguero EG, Geissmann F, Henri S, et al. Constant replenishment from circulating monocytes maintains the macrophage pool in the intestine of adult mice. *Nat Immunol*. 2014;15(10):929-37.
43. Molawi K, Wolf Y, Kandalla PK, Favret J, Hagemeyer N, Frenzel K, et al. Progressive replacement of embryo-derived cardiac macrophages with age. *J Exp Med*. 2014;211(11):2151-8.
44. Murray PJ, Allen JE, Biswas SK, Fisher EA, Gilroy DW, Goerdt S, et al. Macrophage activation and polarization: nomenclature and experimental guidelines. *Immunity*. 2014;41(1):14-20.
45. Laskin DL, Sunil VR, Gardner CR, Laskin JD. Macrophages and tissue injury: agents of defense or destruction? *Annu Rev Pharmacol Toxicol*. 2011;51:267-88.

46. Martinez FO, Gordon S. The M1 and M2 paradigm of macrophage activation: time for reassessment. *F1000Prime Rep.* 2014;6:13.
47. Thompson KK, Tsirka SE. The Diverse Roles of Microglia in the Neurodegenerative Aspects of Central Nervous System (CNS) Autoimmunity. *Int J Mol Sci.* 2017;18(3).
48. Verreck FA, de Boer T, Langenberg DM, Hoeve MA, Kramer M, Vaisberg E, et al. Human IL-23-producing type 1 macrophages promote but IL-10-producing type 2 macrophages subvert immunity to (myco)bacteria. *Proc Natl Acad Sci U S A.* 2004;101(13):4560-5.
49. Kitamura T, Qian BZ, Pollard JW. Immune cell promotion of metastasis. *Nat Rev Immunol.* 2015;15(2):73-86.
50. Rhee I. Diverse macrophages polarization in tumor microenvironment. *Arch Pharm Res.* 2016;39(11):1588-96.
51. Katakura T, Miyazaki M, Kobayashi M, Herndon DN, Suzuki F. CCL17 and IL-10 as effectors that enable alternatively activated macrophages to inhibit the generation of classically activated macrophages. *J Immunol.* 2004;172(3):1407-13.
52. Sica A, Mantovani A. Macrophage plasticity and polarization: in vivo veritas. *J Clin Invest.* 2012;122(3):787-95.
53. Atri C, Guerfali FZ, Laouini D. Role of Human Macrophage Polarization in Inflammation during Infectious Diseases. *Int J Mol Sci.* 2018;19(6).
54. Luo XG, Chen SD. The changing phenotype of microglia from homeostasis to disease. *Transl Neurodegener.* 2012;1(1):9.
55. Tang Y, Le W. Differential Roles of M1 and M2 Microglia in Neurodegenerative Diseases. *Mol Neurobiol.* 2016;53(2):1181-94.
56. Colton C, Wilcock DM. Assessing activation states in microglia. *CNS Neurol Disord Drug Targets.* 2010;9(2):174-91.
57. Block ML, Zecca L, Hong JS. Microglia-mediated neurotoxicity: uncovering the molecular mechanisms. *Nat Rev Neurosci.* 2007;8(1):57-69.
58. Le W, Rowe D, Xie W, Ortiz I, He Y, Appel SH. Microglial activation and dopaminergic cell injury: an in vitro model relevant to Parkinson's disease. *J Neurosci.* 2001;21(21):8447-55.
59. Li R, Huang YG, Fang D, Le WD. (-)-Epigallocatechin gallate inhibits lipopolysaccharide-induced microglial activation and protects against inflammation-mediated dopaminergic neuronal injury. *J Neurosci Res.* 2004;78(5):723-31.
60. Kettenmann H, Hanisch UK, Noda M, Verkhratsky A. Physiology of microglia. *Physiol Rev.* 2011;91(2):461-553.
61. Colton CA. Heterogeneity of microglial activation in the innate immune response in the brain. *J Neuroimmune Pharmacol.* 2009;4(4):399-418.
62. Ponomarev ED, Maresz K, Tan Y, Dittel BN. CNS-derived interleukin-4 is essential for the regulation of autoimmune inflammation and induces a state of alternative activation in microglial cells. *J Neurosci.* 2007;27(40):10714-21.
63. Ledeboer A, Breve JJ, Poole S, Tilders FJ, Van Dam AM. Interleukin-10, interleukin-4, and transforming growth factor-beta differentially regulate lipopolysaccharide-induced production of pro-inflammatory cytokines and nitric oxide in co-cultures of rat astroglial and microglial cells. *Glia.* 2000;30(2):134-42.
64. Suh HS, Zhao ML, Derico L, Choi N, Lee SC. Insulin-like growth factor 1 and 2 (IGF1, IGF2) expression in human microglia: differential regulation by inflammatory mediators. *J Neuroinflammation.* 2013;10:37.
65. Geloso MC, Corvino V, Marchese E, Serrano A, Michetti F, D'Ambrosi N. The Dual Role of Microglia in ALS: Mechanisms and Therapeutic Approaches. *Front Aging Neurosci.* 2017;9:242.

66. Vay SU, Flitsch LJ, Rabenstein M, Rogall R, Blaschke S, Kleinhaus J, et al. The plasticity of primary microglia and their multifaceted effects on endogenous neural stem cells in vitro and in vivo. *J Neuroinflammation*. 2018;15(1):226.
67. Smith TD, Tse MJ, Read EL, Liu WF. Regulation of macrophage polarization and plasticity by complex activation signals. *Integr Biol (Camb)*. 2016;8(9):946-55.
68. Clark CE, Hingorani SR, Mick R, Combs C, Tuveson DA, Vonderheide RH. Dynamics of the immune reaction to pancreatic cancer from inception to invasion. *Cancer Res*. 2007;67(19):9518-27.
69. Diakos CI, Charles KA, McMillan DC, Clarke SJ. Cancer-related inflammation and treatment effectiveness. *Lancet Oncol*. 2014;15(11):e493-503.
70. Candido J, Hagemann T. Cancer-related inflammation. *J Clin Immunol*. 2013;33 Suppl 1:S79-84.
71. Zhang Q, Zhu B, Li Y. Resolution of Cancer-Promoting Inflammation: A New Approach for Anticancer Therapy. *Front Immunol*. 2017;8:71.
72. Nielsen SR, Schmid MC. Macrophages as Key Drivers of Cancer Progression and Metastasis. *Mediators Inflamm*. 2017;2017:9624760.
73. Noy R, Pollard JW. Tumor-associated macrophages: from mechanisms to therapy. *Immunity*. 2014;41(1):49-61.
74. Poon CC, Sarkar S, Yong VW, Kelly JJP. Glioblastoma-associated microglia and macrophages: targets for therapies to improve prognosis. *Brain*. 2017;140(6):1548-60.
75. Chen F, Zhuang X, Lin L, Yu P, Wang Y, Shi Y, et al. New horizons in tumor microenvironment biology: challenges and opportunities. *BMC Med*. 2015;13:45.
76. Wang M, Zhao J, Zhang L, Wei F, Lian Y, Wu Y, et al. Role of tumor microenvironment in tumorigenesis. *J Cancer*. 2017;8(5):761-73.
77. Balkwill FR, Capasso M, Hagemann T. The tumor microenvironment at a glance. *J Cell Sci*. 2012;125(Pt 23):5591-6.
78. Tsai HF, Trubelja A, Shen AQ, Bao G. Tumour-on-a-chip: microfluidic models of tumour morphology, growth and microenvironment. *J R Soc Interface*. 2017;14(131).
79. Dalton HJ, Armaiz-Pena GN, Gonzalez-Villasana V, Lopez-Berestein G, Bar-Eli M, Sood AK. Monocyte subpopulations in angiogenesis. *Cancer Res*. 2014;74(5):1287-93.
80. Silva VL, Al-Jamal WT. Exploiting the cancer niche: Tumor-associated macrophages and hypoxia as promising synergistic targets for nano-based therapy. *J Control Release*. 2017;253:82-96.
81. Komohara Y, Jinushi M, Takeya M. Clinical significance of macrophage heterogeneity in human malignant tumors. *Cancer Sci*. 2014;105(1):1-8.
82. Ostrand-Rosenberg S. Immune surveillance: a balance between protumor and antitumor immunity. *Curr Opin Genet Dev*. 2008;18(1):11-8.
83. Almatroodi SA, McDonald CF, Darby IA, Pouniotis DS. Characterization of M1/M2 Tumour-Associated Macrophages (TAMs) and Th1/Th2 Cytokine Profiles in Patients with NSCLC. *Cancer Microenviron*. 2016;9(1):1-11.
84. Chanmee T, Ontong P, Konno K, Itano N. Tumor-associated macrophages as major players in the tumor microenvironment. *Cancers (Basel)*. 2014;6(3):1670-90.
85. Yang L, Zhang Y. Tumor-associated macrophages: from basic research to clinical application. *J Hematol Oncol*. 2017;10(1):58.
86. Ramanathan S, Jagannathan N. Tumor associated macrophage: a review on the phenotypes, traits and functions. *Iran J Cancer Prev*. 2014;7(1):1-8.
87. Goswami KK, Ghosh T, Ghosh S, Sarkar M, Bose A, Baral R. Tumor promoting role of anti-tumor macrophages in tumor microenvironment. *Cell Immunol*. 2017;316:1-10.
88. Xue J, Schmidt SV, Sander J, Draffehn A, Krebs W, Quester I, et al. Transcriptome-based network analysis reveals a spectrum model of human macrophage activation. *Immunity*. 2014;40(2):274-88.

89. Komohara Y, Horlad H, Ohnishi K, Fujiwara Y, Bai B, Nakagawa T, et al. Importance of direct macrophage-tumor cell interaction on progression of human glioma. *Cancer Sci.* 2012;103(12):2165-72.
90. Komohara Y, Ohnishi K, Kuratsu J, Takeya M. Possible involvement of the M2 anti-inflammatory macrophage phenotype in growth of human gliomas. *J Pathol.* 2008;216(1):15-24.
91. Pyonteck SM, Akkari L, Schuhmacher AJ, Bowman RL, Sevenich L, Quail DF, et al. CSF-1R inhibition alters macrophage polarization and blocks glioma progression. *Nat Med.* 2013;19(10):1264-72.
92. Hambardzumyan D, Gutmann DH, Kettenmann H. The role of microglia and macrophages in glioma maintenance and progression. *Nat Neurosci.* 2016;19(1):20-7.
93. Szulzewsky F, Pelz A, Feng X, Synowitz M, Markovic D, Langmann T, et al. Glioma-associated microglia/macrophages display an expression profile different from M1 and M2 polarization and highly express *Gpnmb* and *Spp1*. *PLoS One.* 2015;10(2):e0116644.
94. Dai XM, Ryan GR, Hapel AJ, Dominguez MG, Russell RG, Kapp S, et al. Targeted disruption of the mouse colony-stimulating factor 1 receptor gene results in osteopetrosis, mononuclear phagocyte deficiency, increased primitive progenitor cell frequencies, and reproductive defects. *Blood.* 2002;99(1):111-20.
95. Mitchell AJ, Roediger B, Weninger W. Monocyte homeostasis and the plasticity of inflammatory monocytes. *Cell Immunol.* 2014;291(1-2):22-31.
96. MacDonald KP, Palmer JS, Cronau S, Seppanen E, Olver S, Raffelt NC, et al. An antibody against the colony-stimulating factor 1 receptor depletes the resident subset of monocytes and tissue- and tumor-associated macrophages but does not inhibit inflammation. *Blood.* 2010;116(19):3955-63.
97. Yona S, Kim KW, Wolf Y, Mildner A, Varol D, Breker M, et al. Fate mapping reveals origins and dynamics of monocytes and tissue macrophages under homeostasis. *Immunity.* 2013;38(1):79-91.
98. Geissmann F, Jung S, Littman DR. Blood monocytes consist of two principal subsets with distinct migratory properties. *Immunity.* 2003;19(1):71-82.
99. Carlin LM, Stamatides EG, Auffray C, Hanna RN, Glover L, Vizcay-Barrena G, et al. Nr4a1-dependent Ly6C(low) monocytes monitor endothelial cells and orchestrate their disposal. *Cell.* 2013;153(2):362-75.
100. Debien E, Mayol K, Biajoux V, Daussy C, De Agüero MG, Taillardet M, et al. S1PR5 is pivotal for the homeostasis of patrolling monocytes. *Eur J Immunol.* 2013;43(6):1667-75.
101. Jakubzick C, Gautier EL, Gibbings SL, Sojka DK, Schlitzer A, Johnson TE, et al. Minimal differentiation of classical monocytes as they survey steady-state tissues and transport antigen to lymph nodes. *Immunity.* 2013;39(3):599-610.
102. Shi C, Jia T, Mendez-Ferrer S, Hohl TM, Serbina NV, Lipuma L, et al. Bone marrow mesenchymal stem and progenitor cells induce monocyte emigration in response to circulating toll-like receptor ligands. *Immunity.* 2011;34(4):590-601.
103. Movahedi K, Laoui D, Gysemans C, Baeten M, Stange G, Van den Bossche J, et al. Different tumor microenvironments contain functionally distinct subsets of macrophages derived from Ly6C(high) monocytes. *Cancer Res.* 2010;70(14):5728-39.
104. Van Overmeire E, Stijlemans B, Heymann F, Keirsse J, Morias Y, Elkrim Y, et al. M-CSF and GM-CSF Receptor Signaling Differentially Regulate Monocyte Maturation and Macrophage Polarization in the Tumor Microenvironment. *Cancer Res.* 2016;76(1):35-42.
105. Scholl SM, Pallud C, Beuvon F, Hacene K, Stanley ER, Rohrschneider L, et al. Anti-colony-stimulating factor-1 antibody staining in primary breast adenocarcinomas correlates with marked inflammatory cell infiltrates and prognosis. *J Natl Cancer Inst.* 1994;86(2):120-6.
106. Lin EY, Nguyen AV, Russell RG, Pollard JW. Colony-stimulating factor 1 promotes progression of mammary tumors to malignancy. *J Exp Med.* 2001;193(6):727-40.
107. Strachan DC, Ruffell B, Oei Y, Bissell MJ, Coussens LM, Pryer N, et al. CSF1R inhibition delays cervical and mammary tumor growth in murine models by attenuating the turnover of tumor-associated macrophages and enhancing infiltration by CD8(+) T cells. *Oncoimmunology.* 2013;2(12):e26968.

108. Platten M, Kretz A, Naumann U, Aulwurm S, Egashira K, Isenmann S, et al. Monocyte chemoattractant protein-1 increases microglial infiltration and aggressiveness of gliomas. *Ann Neurol*. 2003;54(3):388-92.
109. Wu SY, Watabe K. The roles of microglia/macrophages in tumor progression of brain cancer and metastatic disease. *Front Biosci (Landmark Ed)*. 2017;22:1805-29.
110. Ku MC, Wolf SA, Respondek D, Matyash V, Pohlmann A, Waiczies S, et al. GDNF mediates glioblastoma-induced microglia attraction but not astrogliosis. *Acta Neuropathol*. 2013;125(4):609-20.
111. Badie B, Schartner J, Klaver J, Vorpahl J. In vitro modulation of microglia motility by glioma cells is mediated by hepatocyte growth factor/scatter factor. *Neurosurgery*. 1999;44(5):1077-82; discussion 82-3.
112. Epelman S, Lavine KJ, Randolph GJ. Origin and functions of tissue macrophages. *Immunity*. 2014;41(1):21-35.
113. Robbins CS, Hilgendorf I, Weber GF, Theurl I, Iwamoto Y, Figueiredo JL, et al. Local proliferation dominates lesional macrophage accumulation in atherosclerosis. *Nat Med*. 2013;19(9):1166-72.
114. Jenkins SJ, Ruckerl D, Cook PC, Jones LH, Finkelman FD, van Rooijen N, et al. Local macrophage proliferation, rather than recruitment from the blood, is a signature of TH2 inflammation. *Science*. 2011;332(6035):1284-8.
115. Jenkins SJ, Ruckerl D, Thomas GD, Hewitson JP, Duncan S, Brombacher F, et al. IL-4 directly signals tissue-resident macrophages to proliferate beyond homeostatic levels controlled by CSF-1. *J Exp Med*. 2013;210(11):2477-91.
116. Amano SU, Cohen JL, Vangala P, Tencerova M, Nicoloro SM, Yawe JC, et al. Local proliferation of macrophages contributes to obesity-associated adipose tissue inflammation. *Cell Metab*. 2014;19(1):162-71.
117. Tymoszyk P, Evens H, Marzola V, Wachowicz K, Wasmer MH, Datta S, et al. In situ proliferation contributes to accumulation of tumor-associated macrophages in spontaneous mammary tumors. *Eur J Immunol*. 2014;44(8):2247-62.
118. Campbell MJ, Tonlaar NY, Garwood ER, Huo D, Moore DH, Khramtsov AI, et al. Proliferating macrophages associated with high grade, hormone receptor negative breast cancer and poor clinical outcome. *Breast Cancer Res Treat*. 2011;128(3):703-11.
119. Wallerius M, Wallmann T, Bartish M, Ostling J, Mezheyeuski A, Tobin NP, et al. Guidance Molecule SEMA3A Restricts Tumor Growth by Differentially Regulating the Proliferation of Tumor-Associated Macrophages. *Cancer Res*. 2016;76(11):3166-78.
120. Zhu Y, Herndon JM, Sojka DK, Kim KW, Knolhoff BL, Zuo C, et al. Tissue-Resident Macrophages in Pancreatic Ductal Adenocarcinoma Originate from Embryonic Hematopoiesis and Promote Tumor Progression. *Immunity*. 2017;47(2):323-38 e6.
121. Ajami B, Bennett JL, Krieger C, Tetzlaff W, Rossi FM. Local self-renewal can sustain CNS microglia maintenance and function throughout adult life. *Nat Neurosci*. 2007;10(12):1538-43.
122. Bowman RL, Klemm F, Akkari L, Pyonteck SM, Sevenich L, Quail DF, et al. Macrophage Ontogeny Underlies Differences in Tumor-Specific Education in Brain Malignancies. *Cell Rep*. 2016;17(9):2445-59.
123. Giulian D, Ingeman JE. Colony-stimulating factors as promoters of ameboid microglia. *J Neurosci*. 1988;8(12):4707-17.
124. Forstreuter F, Lucius R, Mentlein R. Vascular endothelial growth factor induces chemotaxis and proliferation of microglial cells. *J Neuroimmunol*. 2002;132(1-2):93-8.
125. Coniglio SJ, Eugenin E, Dobrenis K, Stanley ER, West BL, Symons MH, et al. Microglial stimulation of glioblastoma invasion involves epidermal growth factor receptor (EGFR) and colony stimulating factor 1 receptor (CSF-1R) signaling. *Mol Med*. 2012;18:519-27.
126. Arango Duque G, Descoteaux A. Macrophage cytokines: involvement in immunity and infectious diseases. *Front Immunol*. 2014;5:491.
127. Li W, Graeber MB. The molecular profile of microglia under the influence of glioma. *Neuro Oncol*. 2012;14(8):958-78.

128. Hussain SF, Yang D, Suki D, Aldape K, Grimm E, Heimberger AB. The role of human glioma-infiltrating microglia/macrophages in mediating antitumor immune responses. *Neuro Oncol.* 2006;8(3):261-79.
129. De Palma M, Lewis CE. Macrophage regulation of tumor responses to anticancer therapies. *Cancer Cell.* 2013;23(3):277-86.
130. Samaras V, Piperi C, Korkolopoulou P, Zisakis A, Levidou G, Themistocleous MS, et al. Application of the ELISPOT method for comparative analysis of interleukin (IL)-6 and IL-10 secretion in peripheral blood of patients with astroglial tumors. *Mol Cell Biochem.* 2007;304(1-2):343-51.
131. Savage ND, de Boer T, Walburg KV, Joosten SA, van Meijgaarden K, Geluk A, et al. Human anti-inflammatory macrophages induce Foxp3+ GITR+ CD25+ regulatory T cells, which suppress via membrane-bound TGFbeta-1. *J Immunol.* 2008;181(3):2220-6.
132. Cua DJ, Stohlman SA. In vivo effects of T helper cell type 2 cytokines on macrophage antigen-presenting cell induction of T helper subsets. *J Immunol.* 1997;159(12):5834-40.
133. Hanahan D, Folkman J. Patterns and emerging mechanisms of the angiogenic switch during tumorigenesis. *Cell.* 1996;86(3):353-64.
134. Papetti M, Herman IM. Mechanisms of normal and tumor-derived angiogenesis. *Am J Physiol Cell Physiol.* 2002;282(5):C947-70.
135. Bergers G, Benjamin LE. Tumorigenesis and the angiogenic switch. *Nat Rev Cancer.* 2003;3(6):401-10.
136. Folkman J. Tumor angiogenesis: therapeutic implications. *N Engl J Med.* 1971;285(21):1182-6.
137. Baeriswyl V, Christofori G. The angiogenic switch in carcinogenesis. *Semin Cancer Biol.* 2009;19(5):329-37.
138. Relf M, LeJeune S, Scott PA, Fox S, Smith K, Leek R, et al. Expression of the angiogenic factors vascular endothelial cell growth factor, acidic and basic fibroblast growth factor, tumor growth factor beta-1, platelet-derived endothelial cell growth factor, placenta growth factor, and pleiotrophin in human primary breast cancer and its relation to angiogenesis. *Cancer Res.* 1997;57(5):963-9.
139. Kim YW, Byzova TV. Oxidative stress in angiogenesis and vascular disease. *Blood.* 2014;123(5):625-31.
140. Liao D, Johnson RS. Hypoxia: a key regulator of angiogenesis in cancer. *Cancer Metastasis Rev.* 2007;26(2):281-90.
141. Baluk P, Hashizume H, McDonald DM. Cellular abnormalities of blood vessels as targets in cancer. *Curr Opin Genet Dev.* 2005;15(1):102-11.
142. Nagy JA, Chang SH, Shih SC, Dvorak AM, Dvorak HF. Heterogeneity of the tumor vasculature. *Semin Thromb Hemost.* 2010;36(3):321-31.
143. De Palma M, Biziato D, Petrova TV. Microenvironmental regulation of tumour angiogenesis. *Nat Rev Cancer.* 2017;17(8):457-74.
144. Stylianopoulos T, Munn LL, Jain RK. Reengineering the Tumor Vasculature: Improving Drug Delivery and Efficacy. *Trends Cancer.* 2018;4(4):258-9.
145. Coffelt SB, Hughes R, Lewis CE. Tumor-associated macrophages: effectors of angiogenesis and tumor progression. *Biochim Biophys Acta.* 2009;1796(1):11-8.
146. Crowther M, Brown NJ, Bishop ET, Lewis CE. Microenvironmental influence on macrophage regulation of angiogenesis in wounds and malignant tumors. *J Leukoc Biol.* 2001;70(4):478-90.
147. Henze AT, Mazzone M. The impact of hypoxia on tumor-associated macrophages. *J Clin Invest.* 2016;126(10):3672-9.
148. Hagemann T, Robinson SC, Schulz M, Trumper L, Balkwill FR, Binder C. Enhanced invasiveness of breast cancer cell lines upon co-cultivation with macrophages is due to TNF-alpha dependent up-regulation of matrix metalloproteases. *Carcinogenesis.* 2004;25(8):1543-9.
149. Elbarghati L, Murdoch C, Lewis CE. Effects of hypoxia on transcription factor expression in human monocytes and macrophages. *Immunobiology.* 2008;213(9-10):899-908.



150. Fang HY, Hughes R, Murdoch C, Coffelt SB, Biswas SK, Harris AL, et al. Hypoxia-inducible factors 1 and 2 are important transcriptional effectors in primary macrophages experiencing hypoxia. *Blood*. 2009;114(4):844-59.
151. Hildenbrand R, Schaaf A, Dorn-Beineke A, Allgayer H, Sutterlin M, Marx A, et al. Tumor stroma is the predominant uPA-, uPAR-, PAI-1-expressing tissue in human breast cancer: prognostic impact. *Histol Histopathol*. 2009;24(7):869-77.
152. Brandenburg S, Muller A, Turkowski K, Radev YT, Rot S, Schmidt C, et al. Resident microglia rather than peripheral macrophages promote vascularization in brain tumors and are source of alternative pro-angiogenic factors. *Acta Neuropathol*. 2016;131(3):365-78.
153. Hu F, Ku MC, Markovic D, Dzaye OD, Lehnardt S, Synowitz M, et al. Glioma-associated microglial MMP9 expression is upregulated by TLR2 signaling and sensitive to minocycline. *Int J Cancer*. 2014;135(11):2569-78.
154. Giraudo E, Inoue M, Hanahan D. An amino-bisphosphonate targets MMP-9-expressing macrophages and angiogenesis to impair cervical carcinogenesis. *J Clin Invest*. 2004;114(5):623-33.
155. Laoui D, Van Overmeire E, Di Conza G, Aldeni C, Keirsse J, Morias Y, et al. Tumor hypoxia does not drive differentiation of tumor-associated macrophages but rather fine-tunes the M2-like macrophage population. *Cancer Res*. 2014;74(1):24-30.
156. De Palma M, Venneri MA, Galli R, Sergi L, Politi LS, Sampaolesi M, et al. Tie2 identifies a hematopoietic lineage of proangiogenic monocytes required for tumor vessel formation and a mesenchymal population of pericyte progenitors. *Cancer Cell*. 2005;8(3):211-26.
157. Harney AS, Arwert EN, Entenberg D, Wang Y, Guo P, Qian BZ, et al. Real-Time Imaging Reveals Local, Transient Vascular Permeability, and Tumor Cell Intravasation Stimulated by TIE2hi Macrophage-Derived VEGFA. *Cancer Discov*. 2015;5(9):932-43.
158. Matsubara T, Kanto T, Kuroda S, Yoshio S, Higashitani K, Kakita N, et al. TIE2-expressing monocytes as a diagnostic marker for hepatocellular carcinoma correlates with angiogenesis. *Hepatology*. 2013;57(4):1416-25.
159. Oliver G, Srinivasan RS. Lymphatic vasculature development: current concepts. *Ann N Y Acad Sci*. 2008;1131:75-81.
160. Baluk P, Fuxe J, Hashizume H, Romano T, Lashnits E, Butz S, et al. Functionally specialized junctions between endothelial cells of lymphatic vessels. *J Exp Med*. 2007;204(10):2349-62.
161. Stacker SA, Williams SP, Karnezis T, Shayan R, Fox SB, Achen MG. Lymphangiogenesis and lymphatic vessel remodelling in cancer. *Nat Rev Cancer*. 2014;14(3):159-72.
162. Astin JW, Crosier PS. Lymphatics, Cancer and Zebrafish. *Adv Exp Med Biol*. 2016;916:199-218.
163. Semo J, Nicenboim J, Yaniv K. Development of the lymphatic system: new questions and paradigms. *Development*. 2016;143(6):924-35.
164. Jeltsch M, Kaipainen A, Joukov V, Meng X, Lakso M, Rauvala H, et al. Hyperplasia of lymphatic vessels in VEGF-C transgenic mice. *Science*. 1997;276(5317):1423-5.
165. Veikkola T, Jussila L, Makinen T, Karpanen T, Jeltsch M, Petrova TV, et al. Signalling via vascular endothelial growth factor receptor-3 is sufficient for lymphangiogenesis in transgenic mice. *EMBO J*. 2001;20(6):1223-31.
166. Karkkainen MJ, Haiko P, Sainio K, Partanen J, Taipale J, Petrova TV, et al. Vascular endothelial growth factor C is required for sprouting of the first lymphatic vessels from embryonic veins. *Nat Immunol*. 2004;5(1):74-80.
167. Baldwin ME, Halford MM, Roufail S, Williams RA, Hibbs ML, Grail D, et al. Vascular endothelial growth factor D is dispensable for development of the lymphatic system. *Mol Cell Biol*. 2005;25(6):2441-9.
168. Stacker SA, Caesar C, Baldwin ME, Thornton GE, Williams RA, Prevo R, et al. VEGF-D promotes the metastatic spread of tumor cells via the lymphatics. *Nat Med*. 2001;7(2):186-91.

169. Karpanen T, Egeblad M, Karkkainen MJ, Kubo H, Yla-Herttuala S, Jaattela M, et al. Vascular endothelial growth factor C promotes tumor lymphangiogenesis and intralymphatic tumor growth. *Cancer Res.* 2001;61(5):1786-90.
170. Caunt M, Mak J, Liang WC, Stawicki S, Pan Q, Tong RK, et al. Blocking neuropilin-2 function inhibits tumor cell metastasis. *Cancer Cell.* 2008;13(4):331-42.
171. Clasper S, Royston D, Baban D, Cao Y, Ewers S, Butz S, et al. A novel gene expression profile in lymphatics associated with tumor growth and nodal metastasis. *Cancer Res.* 2008;68(18):7293-303.
172. Schoppmann SF, Fenzl A, Nagy K, Unger S, Bayer G, Geleff S, et al. VEGF-C expressing tumor-associated macrophages in lymph node positive breast cancer: impact on lymphangiogenesis and survival. *Surgery.* 2006;139(6):839-46.
173. Wartiovaara U, Salven P, Mikkola H, Lassila R, Kaukonen J, Joukov V, et al. Peripheral blood platelets express VEGF-C and VEGF which are released during platelet activation. *Thromb Haemost.* 1998;80(1):171-5.
174. Schoppmann SF, Birner P, Stockl J, Kalt R, Ullrich R, Caucig C, et al. Tumor-associated macrophages express lymphatic endothelial growth factors and are related to peritumoral lymphangiogenesis. *Am J Pathol.* 2002;161(3):947-56.
175. Skobe M, Hamberg LM, Hawighorst T, Schirner M, Wolf GL, Alitalo K, et al. Concurrent induction of lymphangiogenesis, angiogenesis, and macrophage recruitment by vascular endothelial growth factor-C in melanoma. *Am J Pathol.* 2001;159(3):893-903.
176. Riabov V, Gudima A, Wang N, Mickley A, Orekhov A, Kzhyshkowska J. Role of tumor associated macrophages in tumor angiogenesis and lymphangiogenesis. *Front Physiol.* 2014;5:75.
177. Ding M, Fu X, Tan H, Wang R, Chen Z, Ding S. The effect of vascular endothelial growth factor C expression in tumor-associated macrophages on lymphangiogenesis and lymphatic metastasis in breast cancer. *Mol Med Rep.* 2012;6(5):1023-9.
178. Ji H, Cao R, Yang Y, Zhang Y, Iwamoto H, Lim S, et al. TNFR1 mediates TNF-alpha-induced tumour lymphangiogenesis and metastasis by modulating VEGF-C-VEGFR3 signalling. *Nat Commun.* 2014;5:4944.
179. Solinas G, Germano G, Mantovani A, Allavena P. Tumor-associated macrophages (TAM) as major players of the cancer-related inflammation. *J Leukoc Biol.* 2009;86(5):1065-73.
180. Pollard JW. Tumour-educated macrophages promote tumour progression and metastasis. *Nat Rev Cancer.* 2004;4(1):71-8.
181. Kessenbrock K, Plaks V, Werb Z. Matrix metalloproteinases: regulators of the tumor microenvironment. *Cell.* 2010;141(1):52-67.
182. Mason SD, Joyce JA. Proteolytic networks in cancer. *Trends Cell Biol.* 2011;21(4):228-37.
183. Vempati P, Popel AS, Mac Gabhann F. Extracellular regulation of VEGF: isoforms, proteolysis, and vascular patterning. *Cytokine Growth Factor Rev.* 2014;25(1):1-19.
184. Liu W, Xu J, Wang M, Wang Q, Bi Y, Han M. Tumor-derived vascular endothelial growth factor (VEGF)-a facilitates tumor metastasis through the VEGF-VEGFR1 signaling pathway. *Int J Oncol.* 2011;39(5):1213-20.
185. Mytar B, Woloszyn M, Szatanek R, Baj-Krzyworzeka M, Siedlar M, Ruggiero I, et al. Tumor cell-induced deactivation of human monocytes. *J Leukoc Biol.* 2003;74(6):1094-101.
186. Fonseca AC, Romao L, Amaral RF, Assad Kahn S, Lobo D, Martins S, et al. Microglial stress inducible protein 1 promotes proliferation and migration in human glioblastoma cells. *Neuroscience.* 2012;200:130-41.
187. Lin HC, Song TY, Hu ML. S-Adenosylhomocysteine promotes the invasion of C6 glioma cells via increased secretion of matrix metalloproteinase-2 in murine microglial BV2 cells. *Toxicol Sci.* 2009;112(2):322-30.
188. Goswami S, Sahai E, Wyckoff JB, Cammer M, Cox D, Pixley FJ, et al. Macrophages promote the invasion of breast carcinoma cells via a colony-stimulating factor-1/epidermal growth factor paracrine loop. *Cancer Res.* 2005;65(12):5278-83.

189. Wyckoff JB, Wang Y, Lin EY, Li JF, Goswami S, Stanley ER, et al. Direct visualization of macrophage-assisted tumor cell intravasation in mammary tumors. *Cancer Res.* 2007;67(6):2649-56.
190. Rolny C, Mazzone M, Tugues S, Laoui D, Johansson I, Coulon C, et al. HRG inhibits tumor growth and metastasis by inducing macrophage polarization and vessel normalization through downregulation of PlGF. *Cancer Cell.* 2011;19(1):31-44.
191. Qian BZ, Li J, Zhang H, Kitamura T, Zhang J, Campion LR, et al. CCL2 recruits inflammatory monocytes to facilitate breast-tumour metastasis. *Nature.* 2011;475(7355):222-5.
192. Qian B, Deng Y, Im JH, Muschel RJ, Zou Y, Li J, et al. A distinct macrophage population mediates metastatic breast cancer cell extravasation, establishment and growth. *PLoS One.* 2009;4(8):e6562.
193. Obenauf AC, Massague J. Surviving at a distance: organ specific metastasis. *Trends Cancer.* 2015;1(1):76-91.
194. Hirakawa S, Brown LF, Kodama S, Paavonen K, Alitalo K, Detmar M. VEGF-C-induced lymphangiogenesis in sentinel lymph nodes promotes tumor metastasis to distant sites. *Blood.* 2007;109(3):1010-7.
195. Blackburn HL, Ellsworth DL, Shriver CD, Ellsworth RE. Breast Cancer Metastasis to the Axillary Lymph Nodes: Are Changes to the Lymph Node "Soil" Localized or Systemic? *Breast Cancer (Auckl).* 2017;11:1178223417691246.
196. Ullah I, Karthik GM, Alkodsi A, Kjallquist U, Stalhammar G, Lovrot J, et al. Evolutionary history of metastatic breast cancer reveals minimal seeding from axillary lymph nodes. *J Clin Invest.* 2018;128(4):1355-70.
197. Brown M, Assen FP, Leithner A, Abe J, Schachner H, Asfour G, et al. Lymph node blood vessels provide exit routes for metastatic tumor cell dissemination in mice. *Science.* 2018;359(6382):1408-11.
198. Woo CS, Silberman H, Nakamura SK, Ye W, Sposto R, Colburn W, et al. Lymph node status combined with lymphovascular invasion creates a more powerful tool for predicting outcome in patients with invasive breast cancer. *Am J Surg.* 2002;184(4):337-40.
199. Oberg A, Samii S, Stenling R, Lindmark G. Different occurrence of CD8+, CD45R0+, and CD68+ immune cells in regional lymph node metastases from colorectal cancer as potential prognostic predictors. *Int J Colorectal Dis.* 2002;17(1):25-9.
200. Saijo K, Glass CK. Microglial cell origin and phenotypes in health and disease. *Nat Rev Immunol.* 2011;11(11):775-87.
201. Petty AJ, Yang Y. Tumor-associated macrophages: implications in cancer immunotherapy. *Immunotherapy.* 2017;9(3):289-302.
202. Quail DF, Joyce JA. The Microenvironmental Landscape of Brain Tumors. *Cancer Cell.* 2017;31(3):326-41.
203. Fang H, Declerck YA. Targeting the tumor microenvironment: from understanding pathways to effective clinical trials. *Cancer Res.* 2013;73(16):4965-77.
204. Gazzaniga S, Bravo AI, Guglielmotti A, van Rooijen N, Maschi F, Vecchi A, et al. Targeting tumor-associated macrophages and inhibition of MCP-1 reduce angiogenesis and tumor growth in a human melanoma xenograft. *J Invest Dermatol.* 2007;127(8):2031-41.
205. Dineen SP, Lynn KD, Holloway SE, Miller AF, Sullivan JP, Shames DS, et al. Vascular endothelial growth factor receptor 2 mediates macrophage infiltration into orthotopic pancreatic tumors in mice. *Cancer Res.* 2008;68(11):4340-6.
206. Ries CH, Cannarile MA, Hoves S, Benz J, Wartha K, Runza V, et al. Targeting tumor-associated macrophages with anti-CSF-1R antibody reveals a strategy for cancer therapy. *Cancer Cell.* 2014;25(6):846-59.
207. DeNardo DG, Brennan DJ, Rexhepaj E, Ruffell B, Shiao SL, Madden SF, et al. Leukocyte complexity predicts breast cancer survival and functionally regulates response to chemotherapy. *Cancer Discov.* 2011;1(1):54-67.

208. Lyons YA, Pradeep S, Wu SY, Haemmerle M, Hansen JM, Wagner MJ, et al. Macrophage depletion through colony stimulating factor 1 receptor pathway blockade overcomes adaptive resistance to anti-VEGF therapy. *Oncotarget*. 2017;8(57):96496-505.
209. Xu J, Escamilla J, Mok S, David J, Priceman S, West B, et al. CSF1R signaling blockade stanches tumor-infiltrating myeloid cells and improves the efficacy of radiotherapy in prostate cancer. *Cancer Res*. 2013;73(9):2782-94.
210. Priceman SJ, Sung JL, Shaposhnik Z, Burton JB, Torres-Collado AX, Moughon DL, et al. Targeting distinct tumor-infiltrating myeloid cells by inhibiting CSF-1 receptor: combating tumor evasion of antiangiogenic therapy. *Blood*. 2010;115(7):1461-71.
211. Mok S, Koya RC, Tsui C, Xu J, Robert L, Wu L, et al. Inhibition of CSF-1 receptor improves the antitumor efficacy of adoptive cell transfer immunotherapy. *Cancer Res*. 2014;74(1):153-61.
212. Zhu Y, Knolhoff BL, Meyer MA, Nywening TM, West BL, Luo J, et al. CSF1/CSF1R blockade reprograms tumor-infiltrating macrophages and improves response to T-cell checkpoint immunotherapy in pancreatic cancer models. *Cancer Res*. 2014;74(18):5057-69.
213. Coscia M, Quaglino E, Iezzi M, Curcio C, Pantaleoni F, Riganti C, et al. Zoledronic acid repolarizes tumour-associated macrophages and inhibits mammary carcinogenesis by targeting the mevalonate pathway. *J Cell Mol Med*. 2010;14(12):2803-15.
214. Cieslewicz M, Tang J, Yu JL, Cao H, Zavaljevski M, Motoyama K, et al. Targeted delivery of proapoptotic peptides to tumor-associated macrophages improves survival. *Proc Natl Acad Sci U S A*. 2013;110(40):15919-24.
215. Aslakson CJ, Miller FR. Selective events in the metastatic process defined by analysis of the sequential dissemination of subpopulations of a mouse mammary tumor. *Cancer Res*. 1992;52(6):1399-405.
216. Kaur P, Nagaraja GM, Zheng H, Gizachew D, Galukande M, Krishnan S, et al. A mouse model for triple-negative breast cancer tumor-initiating cells (TNBC-TICs) exhibits similar aggressive phenotype to the human disease. *BMC Cancer*. 2012;12:120.
217. Pulaski BA, Ostrand-Rosenberg S. Mouse 4T1 breast tumor model. *Curr Protoc Immunol*. 2001;Chapter 20:Unit 20 2.
218. Shih AH, Dai C, Hu X, Rosenblum MK, Koutcher JA, Holland EC. Dose-dependent effects of platelet-derived growth factor-B on glial tumorigenesis. *Cancer Res*. 2004;64(14):4783-9.
219. Huse JT, Holland EC. Genetically engineered mouse models of brain cancer and the promise of preclinical testing. *Brain Pathol*. 2009;19(1):132-43.
220. Nakamura F, Kalb RG, Strittmatter SM. Molecular basis of semaphorin-mediated axon guidance. *J Neurobiol*. 2000;44(2):219-29.
221. Kolodkin AL. Semaphorin-mediated neuronal growth cone guidance. *Prog Brain Res*. 1998;117:115-32.
222. Neufeld G, Mumblat Y, Smolkin T, Toledano S, Nir-Zvi I, Ziv K, et al. The role of the semaphorins in cancer. *Cell Adh Migr*. 2016;10(6):652-74.
223. Muller MW, Giese NA, Swiercz JM, Ceyhan GO, Esposito I, Hinz U, et al. Association of axon guidance factor semaphorin 3A with poor outcome in pancreatic cancer. *Int J Cancer*. 2007;121(11):2421-33.
224. Jiang H, Qi L, Wang F, Sun Z, Huang Z, Xi Q. Decreased semaphorin 3A expression is associated with a poor prognosis in patients with epithelial ovarian carcinoma. *Int J Mol Med*. 2015;35(5):1374-80.
225. Tang C, Gao X, Liu H, Jiang T, Zhai X. Decreased expression of SEMA3A is associated with poor prognosis in gastric carcinoma. *Int J Clin Exp Pathol*. 2014;7(8):4782-94.
226. Zhou H, Wu A, Fu W, Lv Z, Zhang Z. Significance of semaphorin-3A and MMP-14 protein expression in non-small cell lung cancer. *Oncol Lett*. 2014;7(5):1395-400.
227. Carrer A, Moimas S, Zacchigna S, Pattarini L, Zentilin L, Ruozi G, et al. Neuropilin-1 identifies a subset of bone marrow Gr1- monocytes that can induce tumor vessel normalization and inhibit tumor growth. *Cancer Res*. 2012;72(24):6371-81.

228. Casazza A, Laoui D, Wenes M, Rizzolio S, Bassani N, Mambretti M, et al. Impeding macrophage entry into hypoxic tumor areas by Sema3A/Nrp1 signaling blockade inhibits angiogenesis and restores antitumor immunity. *Cancer Cell*. 2013;24(6):695-709.
229. Takahashi H, Shibuya M. The vascular endothelial growth factor (VEGF)/VEGF receptor system and its role under physiological and pathological conditions. *Clin Sci (Lond)*. 2005;109(3):227-41.
230. Joukov V, Pajusola K, Kaipainen A, Chilov D, Lahtinen I, Kukk E, et al. A novel vascular endothelial growth factor, VEGF-C, is a ligand for the Flt4 (VEGFR-3) and KDR (VEGFR-2) receptor tyrosine kinases. *EMBO J*. 1996;15(7):1751.
231. Kukk E, Lymboussaki A, Taira S, Kaipainen A, Jeltsch M, Joukov V, et al. VEGF-C receptor binding and pattern of expression with VEGFR-3 suggests a role in lymphatic vascular development. *Development*. 1996;122(12):3829-37.
232. Yuan L, Moyon D, Pardanaud L, Breant C, Karkkainen MJ, Alitalo K, et al. Abnormal lymphatic vessel development in neuropilin 2 mutant mice. *Development*. 2002;129(20):4797-806.
233. Kitsukawa T, Shimono A, Kawakami A, Kondoh H, Fujisawa H. Overexpression of a membrane protein, neuropilin, in chimeric mice causes anomalies in the cardiovascular system, nervous system and limbs. *Development*. 1995;121(12):4309-18.
234. Joukov V, Sorsa T, Kumar V, Jeltsch M, Claesson-Welsh L, Cao Y, et al. Proteolytic processing regulates receptor specificity and activity of VEGF-C. *EMBO J*. 1997;16(13):3898-911.
235. Dixelius J, Makinen T, Wirzenius M, Karkkainen MJ, Wernstedt C, Alitalo K, et al. Ligand-induced vascular endothelial growth factor receptor-3 (VEGFR-3) heterodimerization with VEGFR-2 in primary lymphatic endothelial cells regulates tyrosine phosphorylation sites. *J Biol Chem*. 2003;278(42):40973-9.
236. Nilsson I, Bahram F, Li X, Gualandi L, Koch S, Jarvius M, et al. VEGF receptor 2/-3 heterodimers detected in situ by proximity ligation on angiogenic sprouts. *EMBO J*. 2010;29(8):1377-88.
237. Tammela T, Zarkada G, Nurmi H, Jakobsson L, Heinolainen K, Tvorogov D, et al. VEGFR-3 controls tip to stalk conversion at vessel fusion sites by reinforcing Notch signalling. *Nat Cell Biol*. 2011;13(10):1202-13.
238. Karpanen T, Heckman CA, Keskitalo S, Jeltsch M, Ollila H, Neufeld G, et al. Functional interaction of VEGF-C and VEGF-D with neuropilin receptors. *FASEB J*. 2006;20(9):1462-72.
239. Koch S, Tugues S, Li X, Gualandi L, Claesson-Welsh L. Signal transduction by vascular endothelial growth factor receptors. *Biochem J*. 2011;437(2):169-83.
240. Siegfried G, Basak A, Cromlish JA, Benjannet S, Marcinkiewicz J, Chretien M, et al. The secretory proprotein convertases furin, PC5, and PC7 activate VEGF-C to induce tumorigenesis. *J Clin Invest*. 2003;111(11):1723-32.
241. Su JL, Yang PC, Shih JY, Yang CY, Wei LH, Hsieh CY, et al. The VEGF-C/Flt-4 axis promotes invasion and metastasis of cancer cells. *Cancer Cell*. 2006;9(3):209-23.
242. Shida A, Fujioka S, Kobayashi K, Ishibashi Y, Nimura H, Mitsumori N, et al. Expression of vascular endothelial growth factor (VEGF)-C and -D in gastric carcinoma. *Int J Clin Oncol*. 2006;11(1):38-43.
243. Skobe M, Hawighorst T, Jackson DG, Prevo R, Janes L, Velasco P, et al. Induction of tumor lymphangiogenesis by VEGF-C promotes breast cancer metastasis. *Nat Med*. 2001;7(2):192-8.
244. Mandriota SJ, Jussila L, Jeltsch M, Compagni A, Baetens D, Prevo R, et al. Vascular endothelial growth factor-C-mediated lymphangiogenesis promotes tumour metastasis. *EMBO J*. 2001;20(4):672-82.
245. Valtola R, Salven P, Heikkila P, Taipale J, Joensuu H, Rehn M, et al. VEGFR-3 and its ligand VEGF-C are associated with angiogenesis in breast cancer. *Am J Pathol*. 1999;154(5):1381-90.
246. Moussai D, Mitsui H, Pettersen JS, Pierson KC, Shah KR, Suarez-Farinas M, et al. The human cutaneous squamous cell carcinoma microenvironment is characterized by increased lymphatic density and enhanced expression of macrophage-derived VEGF-C. *J Invest Dermatol*. 2011;131(1):229-36.
247. Zhang B, Wang J, Gao J, Guo Y, Chen X, Wang B, et al. Alternatively activated RAW264.7 macrophages enhance tumor lymphangiogenesis in mouse lung adenocarcinoma. *J Cell Biochem*. 2009;107(1):134-43.

248. Bianco C, Rangel MC, Castro NP, Nagaoka T, Rollman K, Gonzales M, et al. Role of Cripto-1 in stem cell maintenance and malignant progression. *Am J Pathol.* 2010;177(2):532-40.
249. Bianco C, Salomon DS. Targeting the embryonic gene Cripto-1 in cancer and beyond. *Expert Opin Ther Pat.* 2010;20(12):1739-49.
250. Francescangeli F, Contavalli P, De Angelis ML, Baiocchi M, Gambarà G, Pagliuca A, et al. Dynamic regulation of the cancer stem cell compartment by Cripto-1 in colorectal cancer. *Cell Death Differ.* 2015;22(10):1700-13.
251. Phi LTH, Sari IN, Yang YG, Lee SH, Jun N, Kim KS, et al. Cancer Stem Cells (CSCs) in Drug Resistance and their Therapeutic Implications in Cancer Treatment. *Stem Cells Int.* 2018;2018:5416923.
252. Bianco C, Strizzi L, Ebert A, Chang C, Rehman A, Normanno N, et al. Role of human cripto-1 in tumor angiogenesis. *J Natl Cancer Inst.* 2005;97(2):132-41.
253. Rangel MC, Karasawa H, Castro NP, Nagaoka T, Salomon DS, Bianco C. Role of Cripto-1 during epithelial-to-mesenchymal transition in development and cancer. *Am J Pathol.* 2012;180(6):2188-200.
254. Bianco C, Strizzi L, Normanno N, Khan N, Salomon DS. Cripto-1: an oncofetal gene with many faces. *Curr Top Dev Biol.* 2005;67:85-133.
255. Miyoshi N, Ishii H, Mimori K, Sekimoto M, Doki Y, Mori M. TDGF1 is a novel predictive marker for metachronous metastasis of colorectal cancer. *Int J Oncol.* 2010;36(3):563-8.
256. Gong YP, Yarrow PM, Carmalt HL, Kwun SY, Kennedy CW, Lin BP, et al. Overexpression of Cripto and its prognostic significance in breast cancer: a study with long-term survival. *Eur J Surg Oncol.* 2007;33(4):438-43.
257. Ligtenberg MA, Witt K, Galvez-Cancino F, Sette A, Lundqvist A, Lladser A, et al. Cripto-1 vaccination elicits protective immunity against metastatic melanoma. *Oncoimmunology.* 2016;5(5):e1128613.
258. Kazlauskas A. PDGFs and their receptors. *Gene.* 2017;614:1-7.
259. Hermanson M, Funa K, Hartman M, Claesson-Welsh L, Heldin CH, Westermark B, et al. Platelet-derived growth factor and its receptors in human glioma tissue: expression of messenger RNA and protein suggests the presence of autocrine and paracrine loops. *Cancer Res.* 1992;52(11):3213-9.
260. Brennan C, Momota H, Hambardzumyan D, Ozawa T, Tandon A, Pedraza A, et al. Glioblastoma subclasses can be defined by activity among signal transduction pathways and associated genomic alterations. *PLoS One.* 2009;4(11):e7752.
261. Andrae J, Gallini R, Betsholtz C. Role of platelet-derived growth factors in physiology and medicine. *Genes Dev.* 2008;22(10):1276-312.
262. Xian X, Hakansson J, Stahlberg A, Lindblom P, Betsholtz C, Gerhardt H, et al. Pericytes limit tumor cell metastasis. *J Clin Invest.* 2006;116(3):642-51.
263. Di Rocco F, Carroll RS, Zhang J, Black PM. Platelet-derived growth factor and its receptor expression in human oligodendrogliomas. *Neurosurgery.* 1998;42(2):341-6.
264. Kim Y, Kim E, Wu Q, Guryanova O, Hitomi M, Lathia JD, et al. Platelet-derived growth factor receptors differentially inform intertumoral and intratumoral heterogeneity. *Genes Dev.* 2012;26(11):1247-62.